



[Go to Product page](#)

Datasheet for ABIN7159640  
**anti-TET3 antibody (AA 72-335) (FITC)**

### Overview

Quantity:	100 µg
Target:	TET3
Binding Specificity:	AA 72-335
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This TET3 antibody is conjugated to FITC
Application:	Please inquire

### Product Details

Immunogen:	Recombinant Human Methylcytosine dioxygenase TET3 protein (72-335AA)
Isotype:	IgG
Cross-Reactivity:	Human
Purification:	>95%, Protein G purified

### Target Details

Target:	TET3
Alternative Name:	TET3 ( <a href="#">TET3 Products</a> )
Background:	Background: Dioxygenase that catalyzes the conversion of the modified genomic base 5-methylcytosine (5mC) into 5-hydroxymethylcytosine (5hmC) and plays a key role in epigenetic

## Target Details

---

chromatin reprogramming in the zygote following fertilization. Also mediates subsequent conversion of 5hmC into 5-formylcytosine (5fC), and conversion of 5fC to 5-carboxylcytosine (5caC). Conversion of 5mC into 5hmC, 5fC and 5caC probably constitutes the first step in cytosine demethylation. In zygotes, DNA demethylation occurs selectively in the paternal pronucleus before the first cell division, while the adjacent maternal pronucleus and certain paternally-imprinted loci are protected from this process. Participates in DNA demethylation in the paternal pronucleus by mediating conversion of 5mC into 5hmC, 5fC and 5caC. Does not mediate DNA demethylation of maternal pronucleus because of the presence of DPPA3/PGC7 on maternal chromatin that prevents TET3-binding to chromatin (By similarity). In addition to its role in DNA demethylation, also involved in the recruitment of the O-GlcNAc transferase OGT to CpG-rich transcription start sites of active genes, thereby promoting histone H2B GlcNAcylation by OGT.

Aliases: hCG\_40738 antibody, KIAA0401 antibody, Methylcytosine dioxygenase TET3 antibody, Probable methylcytosine dioxygenase TET3 antibody, Tet methylcytosine dioxygenase 3 antibody, Tet oncogene family member 3 antibody, Tet3 antibody, TET3\_HUMAN antibody

---

UniProt: [O43151](#)

---

Pathways: [Warburg Effect](#)

## Application Details

---

Restrictions: For Research Use only

## Handling

---

Format: Liquid

---

Buffer: Preservative: 0.03 % Proclin 300  
Constituents: 50 % Glycerol, 0.01M PBS, PH 7.4

---

Preservative: ProClin

---

Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

---

Storage: -20 °C,-80 °C

---

Storage Comment: Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.