antibodies -online.com







Images



Go to Product page

\sim							
	1//	\Box	$r \setminus$	/ [\bigcirc	1	٨,

Overview		
Quantity:	100 μg	
Target:	MMD2	
Binding Specificity:	AA 1-38	
Reactivity:	Human	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This MMD2 antibody is un-conjugated	
Application:	ELISA, Western Blotting (WB), Immunohistochemistry (IHC), Immunofluorescence (IF)	
Product Details		
Immunogen:	Recombinant Human Monocyte to macrophage differentiation factor 2 protein (1-38AA)	
Isotype:	IgG	
Cross-Reactivity:	Human, Mouse	
Purification:	>95%, Protein G purified	
Target Details		
Target:	MMD2	
Alternative Name:	MMD2 (MMD2 Products)	
Background:	Background: perinuclear region of cytoplasm, protein kinase activity, positive regulation of neuron differentiation, positive regulation of protein kinase activity, positive regulation of Ras	

Target Details

protein signal transduction, protein phosphorylation, regulation of protein localization

Aliases: MMD2 antibody, PAQR10 antibody, Monocyte to macrophage differentiation factor 2

antibody, Progestin and adipoQ receptor family member 10 antibody, Progestin and adipoQ

receptor family member X antibody

UniProt:

Q8IY49


Application Details

Application Notes:	Recommended dilution: WB:1:500-1:2000, IHC:1:20-1:200, IF:1:50-1:200,
Restrictions:	For Research Use only

Handling


папишту		
Format:	Liquid	
Buffer:	Preservative: 0.03 % Proclin 300 Constituents: 50 % Glycerol, 0.01M PBS, PH 7.4	
Preservative:	ProClin	
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.	
Storage:	-20 °C,-80 °C	
Storage Comment:	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.	

Images




Immunohistochemistry

Image 1. Immunohistochemistry of paraffin-embedded human testis tissue using ABIN7160187 at dilution of 1:100



Western Blotting

Image 2. Western blot All lanes: MMD2 antibody at 3 μg/mL Lane 1: Mouse brain tissue Lane 2: Mouse kidney tissue Lane 3: U251 whole cell lysate Lane 4: 293T whole cell lysate Lane 5: Mouse liver tissue Secondary Goat polyclonal to rabbit lgG at 1/10000 dilution Predicted band size: 32, 29, 23 kDa Observed band size: 23 kDa



Immunofluorescence

Image 3. Immunofluorescent analysis of U251 cells using ABIN7160187 at dilution of 1:100 and Alexa Fluor 488-congugated AffiniPure Goat Anti-Rabbit IgG(H+L)