antibodies - online.com







anti-MEF2C antibody (AA 170-380)





Overview

| Quantity: | 100 μL |
|----------------------|--------------------------------------|
| Target: | MEF2C |
| Binding Specificity: | AA 170-380 |
| Reactivity: | Human |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Conjugate: | This MEF2C antibody is un-conjugated |
| Application: | ELISA, Immunohistochemistry (IHC) |

Product Details

| Immunogen: | Recombinant Human Myocyte-specific enhancer factor 2C protein (170-380AA) |
|-------------------|---|
| Isotype: | IgG |
| Cross-Reactivity: | Human |
| Purification: | Antigen Affinity Purified |

Target Details

| Target: | MEF2C |
|-------------------|---|
| Alternative Name: | MEF2C (MEF2C Products) |
| Background: | Background: Transcription activator which binds specifically to the MEF2 element present in |
| | the regulatory regions of many muscle-specific genes. Controls cardiac morphogenesis and |

myogenesis, and is also involved in vascular development. Plays an essential role in hippocampal-dependent learning and memory by suppressing the number of excitatory synapses and thus regulating basal and evoked synaptic transmission. Crucial for normal neuronal development, distribution, and electrical activity in the neocortex. Necessary for proper development of megakaryocytes and platelets and for bone marrow B-lymphopoiesis. Required for B-cell survival and proliferation in response to BCR stimulation, efficient IgG1 antibody responses to T-cell-dependent antigens and for normal induction of germinal center B-cells. May also be involved in neurogenesis and in the development of cortical architecture (By similarity). Isoform 3 and isoform 4, which lack the repressor domain, are more active than isoform 1 and isoform 2.

Aliases: C5DELq14.3 antibody, DEL5q14.3 antibody, MADS box transcription enhancer factor 2 polypeptide C (myocyte enhancer factor 2C) antibody, MADS box transcription enhancer factor 2, polypeptide C antibody, MEF2C antibody, MEF2C_HUMAN antibody, Myocyte enhancer factor 2C antibody, Myocyte specific enhancer factor 2C antibody, Myocyte-specific enhancer factor 2C antibody, OTTHUMP00000222409 antibody, Similar to MADS box transcription enhancer factor 2 polypeptide C antibody

UniProt:

Q06413

Pathways:

Neurotrophin Signaling Pathway, Activation of Innate immune Response, Cellular Response to Molecule of Bacterial Origin, Carbohydrate Homeostasis, Chromatin Binding, Regulation of Muscle Cell Differentiation, Skeletal Muscle Fiber Development, Toll-Like Receptors Cascades, BCR Signaling

Application Details


| Application Notes: | Recommended dilution: IHC:1:20-1:200, |
|--------------------|--|
| Restrictions: | For Research Use only |
| Handling | |
| Format: | Liquid |
| Buffer: | PBS with 0.02 % sodium azide, 50 % glycerol, pH 7.3. |
| Preservative: | Sodium azide |
| Precaution of Use: | This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only. |

Handling

| Storage: | -20 °C,-80 °C |
|----------|---------------|
| | |


Storage Comment: Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.

Images



Immunohistochemistry

Image 1. Immunohistochemistry of paraffin-embedded human colon cancer using ABIN7160504 at dilution of 1:100



Immunohistochemistry

Image 2. Immunohistochemistry of paraffin-embedded human ovarian cancer using ABIN7160504 at dilution of 1:100