antibodies .- online.com







anti-NLRP6 antibody (AA 154-306) (HRP)



()	1/0	r\ /1	014	
()	ve	I V I	-v	V

Quantity:	100 μg
Target:	NLRP6
Binding Specificity:	AA 154-306
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This NLRP6 antibody is conjugated to HRP
Application:	ELISA

Product Details

Immunogen:	Recombinant Human NACHT, LRR and PYD domains-containing protein 6 protein (154-306AA)
Isotype:	IgG
Cross-Reactivity:	Human
Purification:	>95%, Protein G purified

Target Details

Target:	NLRP6
Alternative Name:	NLRP6 (NLRP6 Products)
Background:	Background: As the sensor component of the NLRP6 inflammasome, plays a crucial role in
	innate immunity and inflammation. In response to pathogens and other damage-associated

signals, initiates the formation of the inflammasome polymeric complex, made of NLRP6, PYCARD and CASP1 (and possibly CASP4 and CASP5). Recruitment of proCASP1 to the inflammasome promotes its activation and CASP1-catalyzed IL1B and IL18 maturation and secretion in the extracellular milieu. The precise NLRP6 activation stimulus has not been identified yet (By similarity) (PubMed:12387869). Essential for gut mucosal self-renewal and proliferation. Maintains intestinal homeostasis and a healthy intestinal microbiota. This function is, at least partially, mediated by IL18, and not IL1B, produced by nonhematopoietic cells. Influences intestinal barrier function and microbial homeostasis through the regulation of goblet cell mucus secretion. Acts by promoting autophagy in goblet cells, an essential step for mucus granule exocytosis. Its role in goblet cell physiology is inflammasome-dependent, but IL1B- and IL18-independent. During systemic bacterial infections, may negatively regulate inflammatory signaling and inhibit the influx of monocytes and neutrophils to the circulation and to the peritoneum. May promote peripheral nerve recovery following injury via an inflammasome-independent mechanism (By similarity).

Aliases: Angiotensin II/vasopressin receptor antibody, AVR antibody, CLR11.4 antibody, NACHT, leucine rich repeat and PYD containing 6 antibody, NACHT, LRR and PYD containing protein 6 antibody, NACHT, LRR and PYD domains-containing protein 6 antibody, NAVR antibody, NAVR/AVR antibody, NLR family, pyrin domain containing 6 antibody, Nlrp6 antibody, NLRP6_HUMAN antibody, Nucleotide binding oligomerization domain, leucine rich repeat and pyrin domain containing 6 antibody, PAN3 antibody, PYPAF5 antibody, PYRIN containing APAF1 like protein 5 antibody, PYRIN-containing APAF1-like protein 5 antibody

UniProt: P59044 Pathways:

Inflammasome

Application Details

Application Notes: Optimal working dilution should be determined by the investigator. Restrictions: For Research Use only

Handling

Format: Liquid Buffer: Preservative: 0.03 % Proclin 300 Constituents: 50 % Glycerol, 0.01M PBS, pH 7.4 ProClin Preservative:

Handling

Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.	
Storage:	-20 °C,-80 °C	
Storage Comment:	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.	