

# Datasheet for ABIN7160929 anti-NDUFB6 antibody (AA 2-67)





Go to Product page

_					
	1//	r	Vİ	$\triangle$	۸/
	V		VI		/ V

Quantity:	100 μg	
Target:	NDUFB6	
Binding Specificity:	AA 2-67	
Reactivity:	Human	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This NDUFB6 antibody is un-conjugated	
Application:	ELISA, Western Blotting (WB), Immunoprecipitation (IP), Immunohistochemistry (IHC)	
Product Details		
Immunogen:	Recombinant Human NADH dehydrogenase [ubiquinone] 1 beta subcomplex subunit 6 protein	
Immunogen:	Recombinant Human NADH dehydrogenase [ubiquinone] 1 beta subcomplex subunit 6 protein (2-67AA)	
Immunogen:		
	(2-67AA)	
Isotype:	(2-67AA) IgG	
Isotype: Cross-Reactivity:	(2-67AA)  IgG  Human, Mouse	
Isotype:  Cross-Reactivity:  Purification:	(2-67AA)  IgG  Human, Mouse	
Isotype: Cross-Reactivity: Purification: Target Details	(2-67AA)  IgG  Human, Mouse  >95%, Protein G purified	
Isotype: Cross-Reactivity: Purification: Target Details Target:	(2-67AA)  IgG  Human, Mouse  >95%, Protein G purified  NDUFB6	

#### **Target Details**

dehydrogenase (Complex I), that is believed not to be involved in catalysis. Complex I functions in the transfer of electrons from NADH to the respiratory chain. The immediate electron acceptor for the enzyme is believed to be ubiquinone.

Aliases: B17 antibody, CI antibody, CI B17 antibody, CI-B17 antibody, Complex I mitochondrial respiratory chain B17 subunit antibody, Complex I-B17 antibody, Complex IB17 antibody, MGC13675 antibody, NADH dehydrogenase (ubiquinone) 1 beta subcomplex subunit 6 antibody, NADH dehydrogenase [ubiquinone] 1 beta subcomplex subunit 6 antibody, NADH ubiquinone oxidoreductase B17 subunit antibody, NADH ubiquinone oxidoreductase beta subunit 6 antibody, NADH-ubiquinone oxidoreductase B17 subunit antibody, NDUFB6 antibody

UniProt:

095139

### **Application Details**

Application Notes:	Recommended dilution: WB:1:500-1:5000, IHC:1:20-1:200, IP:1:200-1:2000,	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Buffer:	Preservative: 0.03 % Proclin 300	
	Constituents: 50 % Glycerol, 0.01M PBS, PH 7.4	
Preservative:	ProClin	


Constituents: 50 % Glycerol, 0.01M PBS, PH 7.4

Preservative: ProClin

Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.


Storage: -20 °C,-80 °C

Storage Comment: Upon receipt, store at -20 °C or -80 °C. Avoid repeated freeze.




#### **Immunohistochemistry**

**Image 1.** Immunohistochemistry of paraffin-embedded human lung cancer using ABIN7160929 at dilution of 1:100



## **Western Blotting**

Image 2. Western Blot Positive WB detected in: HEK293 whole cell lysate, Mouse heart tissue, Mouse liver tissue, Mouse spleen tissue, Mouse lung tissue, Mouse kidney tissue, Mouse brain tissue, Mouse skeletal muscle tissue, Mouse thymus tissue All lanes: NDUFB6 antibody at 3  $\mu$  g/mL Secondary Goat polyclonal to rabbit IgG at 1/50000 dilution Predicted band size: 16, 14 kDa Observed band size: 16, 14 kDa



#### **Western Blotting**

**Image 3.** Immunoprecipitating NDUFB6 in HEK293 whole cell lysate Lane 1: Rabbit control IgG instead of (1  $\mu$ g) instead of ABIN7160929 in HEK293 whole cell lysate. For western blotting, a HRP-conjugated Protein G antibody was used as the secondary antibody (1/2000) Lane 2: ABIN7160929 (4  $\mu$ g) + HEK293 whole cell lysate (500  $\mu$ g) Lane 3: HEK293 whole cell lysate (20  $\mu$ g)