## antibodies -online.com





## anti-NMNAT3 antibody (AA 1-215) (Biotin)



( )	1/0	r\ /1	014	
( )	ve	I V I	-v	V

Quantity:	100 μg	
Target:	NMNAT3	
Binding Specificity:	AA 1-215	
Reactivity:	Human	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This NMNAT3 antibody is conjugated to Biotin	
Application:	ELISA	
Product Details		
Immunogen:	Recombinant Human Nicotinamide/nicotinic acid mononucleotide adenylyltransferase 3 protein (1-215AA)	
Isotype:	IgG	
Cross-Reactivity:	Human	
Purification:	>95%, Protein G purified	
Target Details		
Target:	NMNAT3	
Alternative Name:	NMNAT3 (NMNAT3 Products)	
Background:	Background: Catalyzes the formation of NAD+ from nicotinamide mononucleotide (NMN) and	

ATP. Can also use the deamidated form, nicotinic acid mononucleotide (NaMN) as substrate with the same efficiency. Can use triazofurin monophosphate (TrMP) as substrate. Can also use GTP and ITP as nucleotide donors. Also catalyzes the reverse reaction, i.e. the pyrophosphorolytic cleavage of NAD+. For the pyrophosphorolytic activity, can use NAD (+), NADH, NAAD, nicotinic acid adenine dinucleotide phosphate (NHD), nicotinamide guanine dinucleotide (NGD) as substrates. Fails to cleave phosphorylated dinucleotides NADP+, NADPH and NAADP+. Protects against axonal degeneration following injury.

Aliases: NaMN adenylyltransferase 3 antibody, Nicotinamide mononucleotide adenylyltransferase 3 antibody, Nicotinamide nucleotide adenylyltransferase 3 antibody, Nicotinate-nucleotide adenylyltransferase 3 antibody, NMN adenylyltransferase 3 antibody, NMNA3\_HUMAN antibody, NMNAT 3 antibody, Nmnat3 antibody, PNAT 3 antibody, PNAT-3 antibody, Pyridine nucleotide adenylyltransferase 3 antibody

UniProt:

Q96T66

## **Application Details**

Application Notes:	Optimal working dilution should be determined by the investigator.	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Buffer:	Preservative: 0.03 % Proclin 300 Constituents: 50 % Glycerol, 0.01M PBS, PH 7.4	
Preservative:	ProClin	
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.	
Storage:	-20 °C,-80 °C	
Storage Comment:	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.	