

Datasheet for ABIN7161800

anti-NFATC3 antibody (AA 887-1070) (HRP)[Go to Product page](#)

Overview

Quantity:	100 µg
Target:	NFATC3
Binding Specificity:	AA 887-1070
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This NFATC3 antibody is conjugated to HRP
Application:	ELISA

Product Details

Immunogen:	Recombinant Human Nuclear factor of activated T-cells, cytoplasmic 3 protein (887-1070AA)
Isotype:	IgG
Cross-Reactivity:	Human
Purification:	>95%, Protein G purified

Target Details

Target:	NFATC3
Alternative Name:	NFATC3 (NFATC3 Products)
Background:	Background: Acts as a regulator of transcriptional activation. Plays a role in the inducible expression of cytokine genes in T-cells, especially in the induction of the IL-2.

Target Details

Aliases: C80703 antibody, cytoplasmic 3 antibody, D8ErtD281e antibody, NF AT4 antibody, NF ATc3 antibody, NF-AT4 antibody, NF-ATc3 antibody, NFAC3_HUMAN antibody, NFAT4 antibody, NFATc3 antibody, NFATx antibody, Nuclear factor of activated T cells, cytoplasmic 3 antibody, nuclear factor of activated T cells, cytoplasmic, calcineurin dependent 3 antibody, Nuclear factor of activated T-cells antibody, T cell transcription factor NFAT4 antibody, T-cell transcription factor NFAT4 antibody

UniProt:	Q12968
Pathways:	RTK Signaling , WNT Signaling , Fc-epsilon Receptor Signaling Pathway , Chromatin Binding

Application Details

Application Notes:	Optimal working dilution should be determined by the investigator.
Restrictions:	For Research Use only

Handling

Format:	Liquid
Buffer:	Preservative: 0.03 % Proclin 300 Constituents: 50 % Glycerol, 0.01M PBS, pH 7.4
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C,-80 °C
Storage Comment:	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.