antibodies .- online.com







anti-PNLIP antibody (AA 60-214)



Image



Overview

| Quantity: | 100 μg |
|----------------------|--------------------------------------|
| Target: | PNLIP |
| Binding Specificity: | AA 60-214 |
| Reactivity: | Human |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Conjugate: | This PNLIP antibody is un-conjugated |
| Application: | ELISA, Immunohistochemistry (IHC) |

Product Details

| Immunogen: | Recombinant Human Pancreatic triacylglycerol lipase protein (60-214AA) |
|-------------------|--|
| Isotype: | IgG |
| Cross-Reactivity: | Human |
| Purification: | >95%, Protein G purified |

Target Details

| Target: | PNLIP |
|-------------------|--|
| Alternative Name: | PNLIP (PNLIP Products) |
| Background: | Background: extracellular region, triglyceride lipase activity, lipid digestion, positive regulation of triglyceride lipase activity, retinoid metabolic process |

Target Details

| Aliases: lipase, pancreatic antibody, LIPP_HUMAN antibody, Pancreatic lipase | antibody, |
|--|-----------------|
| Pancreatic triacylglycerol lipase antibody, PL antibody, PNLIP antibody, PNLIP | D antibody, PTL |
| antibody, Triacylglycerol acylhydrolase antibody | |
| | |

UniProt:

P16233


Application Details

| Application Notes: | Recommended dilution: IHC:1:200-1:500, |
|--------------------|--|
| Restrictions: | For Research Use only |

Handling

| Format: | Liquid |
|--------------------|---|
| Buffer: | Preservative: 0.03 % Proclin 300 Constituents: 50 % Glycerol, 0.01M PBS, pH 7.4 |
| Preservative: | ProClin |
| Precaution of Use: | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only. |
| Storage: | -20 °C,-80 °C |
| Storage Comment: | Upon receipt, store at -20°C or -80°C. Avoid repeated freeze. |

Images



Immunohistochemistry

Image 1. IHC image of ABIN7162579 diluted at 1:200 and staining in paraffin-embedded human pancreatic tissue performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.