antibodies -online.com





anti-WASF2 antibody (AA 131-230) (Biotin)



Overview

Quantity:	100 μL
Target:	WASF2
Binding Specificity:	AA 131-230
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This WASF2 antibody is conjugated to Biotin
Application:	ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human WASF2
Isotype:	IgG
Predicted Reactivity:	Human,Mouse,Rat,Dog,Cow,Sheep,Chicken
Purification:	Purified by Protein A.

Target Details

Target:	WASF2
Alternative Name:	WASF2 (WASF2 Products)
Background:	Synonyms: Protein WAVE-2, SCAR 2, SCAR, SCAR2, Verprolin homology domain-containing

protein 2, WASF2, WASF2_HUMAN, WASP family protein member 2, WASP-family protein member 2, WAVE 2, WAVE2, Wiskott-Aldrich syndrome protein family member 2, Wiskott-Aldrich syndrome protein family verprolin-homologous protein, DICTYOSTELIUM, WASF2_HUMAN.

Background: This gene encodes a member of the Wiskott-Aldrich syndrome protein family. The gene product is a protein that forms a multiprotein complex that links receptor kinases and actin. Binding to actin occurs through a C-terminal verprolin homology domain in all family members. The multiprotein complex serves to tranduce signals that involve changes in cell shape, motility or function. The published map location (PMID:10381382) has been changed based on recent genomic sequence comparisons, which indicate that the expressed gene is located on chromosome 1, and a pseudogene may be located on chromosome X. Two transcript variants encoding different isoforms have been found for this gene.

Gene ID: 10163

Pathways: RTK Signaling

Application Details

Application Notes: IHC-P 1:200-400

IHC-F 1:100-500

Restrictions: For Research Use only

Handling

Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C for 12 months.
Expiry Date:	12 months