

Datasheet for ABIN7162790
anti-PPIA antibody (AA 1-165)



[Go to Product page](#)

1 Image

Overview

Quantity:	100 µg
Target:	PPIA
Binding Specificity:	AA 1-165
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This PPIA antibody is un-conjugated
Application:	Western Blotting (WB), ELISA

Product Details

Immunogen:	Recombinant Human Peptidyl-prolyl cis-trans isomerase A protein (1-165AA)
Isotype:	IgG
Cross-Reactivity:	Human, Mouse
Purification:	>95%, Protein G purified

Target Details

Target:	PPIA
Alternative Name:	PPIA (PPIA Products)
Background:	Background: PPIases accelerate the folding of proteins. It catalyzes the cis-trans isomerization of proline imidic peptide bonds in oligopeptides.

Target Details

Aliases: Cyclophilin A antibody, Cyclophilin antibody, CyclophilinA antibody, Cyclosporin A binding protein antibody, Cyclosporin A-binding protein antibody, CYPA antibody, CYPH antibody, Epididymis secretory sperm binding protein Li 69p antibody, HEL S 69p antibody, MGC117158 antibody, MGC12404 antibody, MGC23397 antibody, Peptidyl prolyl cis trans isomerase A antibody, Peptidyl-prolyl cis-trans isomerase A antibody, Peptidylprolyl isomerase A (cyclophilin A) antibody, Peptidylprolyl isomerase A antibody, PPIA antibody, PPIA protein antibody, PPIA_HUMAN antibody, PPIase A antibody, Rotamase A antibody, RotamaseA antibody, T cell cyclophilin antibody

UniProt: [P62937](#)

Application Details

Application Notes: Recommended dilution: WB:1:1000-1:5000,

Restrictions: For Research Use only

Handling

Format: Liquid


Buffer: Preservative: 0.03 % Proclin 300
Constituents: 50 % Glycerol, 0.01M PBS, PH 7.4

Preservative: ProClin

Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Storage: -20 °C,-80 °C

Storage Comment: Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.



Western Blotting

Image 1. Western Blot Positive WB detected in: HepG2 whole cell lysate, Mouse kidney tissue, Mouse brain tissue
All lanes: PPIA antibody at 4 µg/mL Secondary Goat polyclonal to rabbit IgG at 1/50000 dilution Predicted band size: 19, 12 kDa Observed band size: 19 kDa