

Datasheet for ABIN7163060
anti-PHF7 antibody (AA 277-368)[Go to Product page](#)

2 Images

Overview

Quantity:	100 µg
Target:	PHF7
Binding Specificity:	AA 277-368
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This PHF7 antibody is un-conjugated
Application:	ELISA, Immunohistochemistry (IHC)

Product Details

Immunogen:	Recombinant Human PHD finger protein 7 protein (277-368AA)
Isotype:	IgG
Cross-Reactivity:	Human
Purification:	>95%, Protein G purified

Target Details

Target:	PHF7
Alternative Name:	PHF7 (PHF7 Products)
Background:	Background: May play a role in spermatogenesis. Aliases: DKFZp434L1850 antibody, HSPC045 antibody, HSPC226 antibody, MGC26088

Target Details

antibody, NYD SP6 antibody, OTTHUMP00000212988 antibody, OTTHUMP00000212989 antibody, OTTHUMP00000212990 antibody, OTTHUMP00000212993 antibody, PHD finger protein 7 antibody, Phf7 antibody, PHF7_HUMAN antibody, Testis development protein NYD SP6 antibody, Testis development protein NYD-SP6 antibody

UniProt: [Q9BW1](#)

Application Details

Application Notes: Recommended dilution: IHC:1:200-1:500,

Restrictions: For Research Use only

Handling

Format: Liquid

Buffer: Preservative: 0.03 % Proclin 300
Constituents: 50 % Glycerol, 0.01M PBS, pH 7.4


Preservative: ProClin

Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Storage: -20 °C,-80 °C


Storage Comment: Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.

Images



Immunohistochemistry

Image 1. IHC image of ABIN7163060 diluted at 1:300 and staining in paraffin-embedded human endometrial cancer performed on a Leica Bond™ system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10 % normal goat serum 30 min at RT. Then primary antibody (1 % BSA) was incubated at 4 °C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.



Immunohistochemistry

Image 2. IHC image of ABIN7163060 diluted at 1:300 and staining in paraffin-embedded human testis tissue performed on a Leica Bond™ system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10 % normal goat serum 30 min at RT. Then primary antibody (1 % BSA) was incubated at 4 °C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.