antibodies -online.com





anti-PREX2 antibody (AA 1417-1592) (HRP)



Go to Product page

| \sim | | | | | |
|--------|-----|----|-----|-------------|----|
| | 1/6 | ⊃r | \/I | \triangle | ٨/ |

Alternative Name:

Background:

| Quantity: | 100 μg | |
|----------------------|--|--|
| Target: | PREX2 (DEPDC2) | |
| Binding Specificity: | AA 1417-1592 | |
| Reactivity: | Human | |
| Host: | Rabbit | |
| Clonality: | Polyclonal | |
| Conjugate: | This PREX2 antibody is conjugated to HRP | |
| Application: | ELISA | |
| Product Details | | |
| Immunogen: | Recombinant Human Phosphatidylinositol 3,4,5-trisphosphate-dependent Rac exchanger 2 | |
| | protein (1417-1592AA) | |
| Isotype: | IgG | |
| Cross-Reactivity: | Human | |
| Purification: | >95%, Protein G purified | |
| Target Details | | |
| Target: | PREX2 (DEPDC2) | |

Background: Functions as a RAC1 guanine nucleotide exchange factor (GEF), activating Rac

PREX2 (DEPDC2 Products)

proteins by exchanging bound GDP for free GTP. Its activity is synergistically activated by phosphatidylinositol 3,4,5-trisphosphate and the beta gamma subunits of heterotrimeric G protein. Mediates the activation of RAC1 in a PI3K-dependent manner. May be an important mediator of Rac signaling, acting directly downstream of both G protein-coupled receptors and phosphoinositide 3-kinase.

Aliases: 4 antibody, 5)-dependent Rac exchanger 2 antibody, 5-trisphosphate-dependent Rac exchanger 2 protein antibody, 6230420N16Rik antibody, DEP domain-containing protein 2 antibody, DEP.2 antibody, DEPDC2 antibody, FLJ12987 antibody, P-Rex2 antibody, Phosphatidylinositol 3 antibody, phosphatidylinositol 3,4,5-trisphosphate-dependent Rac exchanger 2 antibody, phosphatidylinositol 3,4,5-trisphosphate-dependent Rac exchanger 2 protein antibody, phosphatidylinositol-3,4,5-trisphosphate-dependent Rac exchange factor 2 antibody, PREX2 antibody, PREX2_HUMAN antibody, PtdIns(3 antibody, PtdIns(3,4,5)-dependent Rac exchanger 2 antibody

UniProt:

Q70Z35

Application Details

| Application Notes: | Optimal working dilution should be determined by the investigator. | |
|--------------------|---|--|
| Restrictions: | For Research Use only | |
| Handling | | |
| Format: | Liquid | |
| Buffer: | Preservative: 0.03 % Proclin 300 Constituents: 50 % Glycerol, 0.01M PBS, PH 7.4 | |
| Preservative: | ProClin | |
| Precaution of Use: | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only. | |
| Storage: | -20 °C,-80 °C | |
| Storage Comment: | Upon receipt, store at -20°C or -80°C. Avoid repeated freeze. | |