

Datasheet for ABIN7163189

anti-PIK3CA antibody (Catalytic Subunit alpha) (Biotin)



Go to Product page

Overview	
Quantity:	100 μg
Target:	PIK3CA
Binding Specificity:	AA 1-206, Catalytic Subunit alpha
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This PIK3CA antibody is conjugated to Biotin
Application:	ELISA
Product Details	
Immunogen:	Recombinant Human Phosphatidylinositol 4,5-bisphosphate 3-kinase catalytic subunit alpha
	isoform protein (1-206AA)
Isotype:	lgG
Cross-Reactivity:	Human
Purification:	>95%, Protein G purified
Target Details	
Target:	PIK3CA
Alternative Name:	PIK3CA (PIK3CA Products)
Background:	Background: Phosphoinositide-3-kinase (PI3K) that phosphorylates PtdIns

(Phosphatidylinositol), PtdIns4P (Phosphatidylinositol 4-phosphate) and PtdIns(4,5)P2 (Phosphatidylinositol 4,5-bisphosphate) to generate phosphatidylinositol 3,4,5-trisphosphate (PIP3). PIP3 plays a key role by recruiting PH domain-containing proteins to the membrane, including AKT1 and PDPK1, activating signaling cascades involved in cell growth, survival, proliferation, motility and morphology. Participates in cellular signaling in response to various growth factors. Involved in the activation of AKT1 upon stimulation by receptor tyrosine kinases ligands such as EGF, insulin, IGF1, VEGFA and PDGF. Involved in signaling via insulin-receptor substrate (IRS) proteins. Essential in endothelial cell migration during vascular development through VEGFA signaling, possibly by regulating RhoA activity. Required for lymphatic vasculature development, possibly by binding to RAS and by activation by EGF and FGF2, but not by PDGF. Regulates invadopodia formation through the PDPK1-AKT1 pathway. Participates in cardiomyogenesis in embryonic stem cells through a AKT1 pathway. Participates in vasculogenesis in embryonic stem cells through PDK1 and protein kinase C pathway. Also has serine-protein kinase activity: phosphorylates PIK3R1 (p85alpha regulatory subunit), EIF4EBP1 and HRAS. Plays a role in the positive regulation of phagocytosis and pinocytosis (By similarity). Aliases: 5-bisphosphate 3-kinase 110 kDa catalytic subunit alpha antibody, 5-bisphosphate 3kinase catalytic subunit alpha isoform antibody, caPI3K antibody, CLOVE antibody, CWS5 antibody, MCAP antibody, MCM antibody, MCMTC antibody, MGC142161 antibody, MGC142163 antibody, p110 alpha antibody, p110alpha antibody, Phosphatidylinositol 3 kinase catalytic alpha polypeptide antibody, Phosphatidylinositol 3 kinase catalytic 110 KD alpha antibody, Phosphatidylinositol 4 5 bisphosphate 3 kinase catalytic subunit alpha antibody, Phosphatidylinositol 4 5 bisphosphate 3 kinase catalytic subunit alpha isoform antibody, Phosphatidylinositol 4,5 bisphosphate 3 kinase 110 kDa catalytic subunit alpha antibody, Phosphatidylinositol-4 antibody, Phosphoinositide 3 kinase catalytic alpha polypeptide antibody, PI 3 Kinase catalytic subunit alpha antibody, PI3 kinase p110 subunit alpha antibody, PI3-kinase subunit alpha antibody, PI3K antibody, PI3K-alpha antibody, PI3KC A antibody, PIK3C A antibody, Pik3ca antibody, PK3CA antibody, PK3CA_HUMAN antibody, PtdIns 3 kinase p110 antibody, Ptdlns-3-kinase subunit alpha antibody, Ptdlns-3-kinase subunit p110-alpha antibody, Serine/threonine protein kinase PIK3CA antibody

UniProt:

P42336

Pathways:

PI3K-Akt Signaling, RTK Signaling, TCR Signaling, AMPK Signaling, Interferon-gamma Pathway, TLR Signaling, Fc-epsilon Receptor Signaling Pathway, EGFR Signaling Pathway, Neurotrophin Signaling Pathway, Inositol Metabolic Process, Hepatitis C, CXCR4-mediated Signaling Events, Signaling Events mediated by VEGFR1 and VEGFR2, Signaling of Hepatocyte Growth Factor Receptor, VEGFR1 Specific Signals, VEGF Signaling

Application Details

Application Notes:	Optimal working dilution should be determined by the investigator.
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Buffer:	Preservative: 0.03 % Proclin 300 Constituents: 50 % Glycerol, 0.01M PBS, pH 7.4
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C,-80 °C
Storage Comment:	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.