antibodies -online.com





anti-ATP8B1 antibody (AA 217-330) (Biotin)



| () | ve | K\ / | | |
|-----|--------|------|---|----|
| | \cup | 1 V/ | - | V۷ |
| | | | | |

| Quantity: | 100 μg |
|----------------------|--|
| Target: | ATP8B1 |
| Binding Specificity: | AA 217-330 |
| Reactivity: | Human |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Conjugate: | This ATP8B1 antibody is conjugated to Biotin |
| Application: | ELISA |

Product Details

| Immunogen: | Recombinant Human Phospholipid-transporting ATPase IC protein (217-330AA) | |
|-------------------|---|--|
| Isotype: | IgG | |
| Cross-Reactivity: | Human | |
| Purification: | >95%, Protein G purified | |

Target Details

| Target: | ATP8B1 | |
|-------------------|--|--|
| Alternative Name: | ATP8B1 (ATP8B1 Products) | |
| Background: | Background: Catalytic component of a P4-ATPase flippase complex which catalyzes the | |
| | hydrolysis of ATP coupled to the transport of aminophospholipids from the outer to the inner | |

leaflet of various membranes and ensures the maintenance of asymmetric distribution of phospholipids. Phospholipid translocation seems also to be implicated in vesicle formation and in uptake of lipid signaling molecules. May play a role in asymmetric distribution of phospholipids in the canicular membrane. May have a role in transport of bile acids into the canaliculus, uptake of bile acids from intestinal contents into intestinal mucosa or both. In cooperation with ABCB4 may be involved in establishing integrity of the canalicular membrane thus protecting hepatocytes from bile salts. Together with TMEM30A is involved in uptake of the synthetic drug alkylphospholipid perifosine. Involved in the microvillus formation in polarized epithelial cells, the function seems to be independent from its flippase activity. Required for the preservation of cochlear hair cells in the inner ear. May act as cardiolipin transporter during inflammatory injury.

Aliases: AT8B1_HUMAN antibody, ATP8B1 antibody, ATPase class I type 8B member 1 antibody, ATPase, aminophospholipid transporter, class I, type 8B, member 1 antibody, ATPIC antibody, BRIC antibody, E1-E2 ATPase antibody, Familial intrahepatic cholestasis type 1 antibody, FIC1 antibody, OTTHUMP00000163615 antibody, PFIC antibody, PFIC1 antibody, Phospholipid transporting ATPase IC antibody, Probable phospholipid-transporting ATPase IC antibody

UniProt:

043520

Application Details

| Application Notes: | Optimal working dilution should be determined by the investigator. |
|--------------------|--|
| Restrictions: | For Research Use only |

Handling

| Format: | Liquid | |
|--------------------|---|--|
| | | |
| Buffer: | Preservative: 0.03 % Proclin 300 | |
| | Constituents: 50 % Glycerol, 0.01M PBS, PH 7.4 | |
| Preservative: | ProClin | |
| Precaution of Use: | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only. | |
| Storage: | -20 °C,-80 °C | |
| Storage Comment: | Upon receipt, store at -20°C or -80°C. Avoid repeated freeze. | |