antibodies -online.com





anti-PITX3 antibody (AA 10-200)

3 Images



Overview

Quantity:	100 μg
Target:	PITX3
Binding Specificity:	AA 10-200
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This PITX3 antibody is un-conjugated
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (IHC)

Product Details

Immunogen:	Recombinant Human Pituitary homeobox 3 protein (10-200AA)
Isotype:	IgG
Cross-Reactivity:	Human, Mouse
Purification:	>95%, Protein G purified

Target Details

Target:	PITX3
Alternative Name:	PITX3 (PITX3 Products)
Background:	Background: Transcriptional regulator which is important for the differentiation and
	maintenance of meso-diencephalic dopaminergic (mdDA) neurons during development. In

addition to its importance during development, it also has roles in the long-term survival and maintenance of the mdDA neurons. Activates NR4A2/NURR1-mediated transcription of genes such as SLC6A3, SLC18A2, TH and DRD2 which are essential for development of mdDA neurons. Acts by decreasing the interaction of NR4A2/NURR1 with the corepressor NCOR2/SMRT which acts through histone deacetylases (HDACs) to keep promoters of NR4A2/NURR1 target genes in a repressed deacetylated state. Essential for the normal lens development and differentiation. Plays a critical role in the maintenance of mitotic activity of lens epithelial cells, fiber cell differentiation and in the control of the temporal and spatial activation of fiber cell-specific crystallins. Positively regulates FOXE3 expression and negatively regulates PROX1 in the anterior lens epithelium, preventing activation of CDKN1B/P27Kip1 and CDKN1C/P57Kip2 and thus maintains lens epithelial cells in cell cycle (By similarity).

Aliases: Homeobox protein PITX 3 antibody, Homeobox protein PITX3 antibody, MGC12766 antibody, Paired like homeodomain transcription factor 3 antibody, Pitx3 antibody, Pitx3 antibody, PITX3_HUMAN antibody, PTX 3 antibody, PTX3 antibody

UniProt:


075364

Application Details

Application Notes:


Application Notes:	Recommended dilution: WB:1:1000-1:5000, IHC:1:20-1:200,
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Buffer:	Preservative: 0.03 % Proclin 300 Constituents: 50 % Glycerol, 0.01M PBS, PH 7.4
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C,-80 °C
Storage Comment:	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.

Pagement and dilution: WP:1:1000 1:5000 IHC:1:20 1:200




Immunohistochemistry

Image 1. Immunohistochemistry of paraffin-embedded human kidney tissue using ABIN7163469 at dilution of 1:100



Western Blotting

Image 2. Western Blot Positive WB detected in: Mouse spleen tissue, Mouse lung tissue, Mouse brain tissue All lanes: PITX3 antibody at 2.7 µg/mL Secondary Goat polyclonal to rabbit IgG at 1/50000 dilution Predicted band size: 32 kDa Observed band size: 32 kDa



Immunohistochemistry

Image 3. Immunohistochemistry of paraffin-embedded human placenta tissue using ABIN7163469 at dilution of 1:100