antibodies - online.com







anti-PIWIL4 antibody (AA 12-235)





Overview

Quantity:	100 μg
Target:	PIWIL4
Binding Specificity:	AA 12-235
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This PIWIL4 antibody is un-conjugated
Application:	Western Blotting (WB), ELISA, Immunofluorescence (IF)

Product Details

Immunogen:	Recombinant Human Piwi-like protein 4 protein (12-235AA)
Isotype:	IgG
Cross-Reactivity:	Human
Purification:	>95%, Protein G purified

Target Details

Target:	PIWIL4
Alternative Name:	PIWIL4 (PIWIL4 Products)
Background:	Background: Plays a central role during spermatogenesis by repressing transposable elements
	and preventing their mobilization, which is essential for the germline integrity. Acts via the

piRNA metabolic process, which mediates the repression of transposable elements during meiosis by forming complexes composed of piRNAs and Piwi proteins and governs the methylation and subsequent repression of transposons. Directly binds piRNAs, a class of 24 to 30 nucleotide RNAs that are generated by a Dicer-independent mechanism and are primarily derived from transposons and other repeated sequence elements. Associates with secondary piRNAs antisense and PIWIL2/MILI is required for such association. The piRNA process acts upstream of known mediators of DNA methylation. Participates in a piRNA amplification loop. Besides their function in transposable elements repression, piRNAs are probably involved in other processes during meiosis such as translation regulation (By similarity). May be involved in the chromatin-modifying pathway by inducing \\\'Lys-9\\\' methylation of histone H3 at some loci (PubMed:12906857).

Aliases: DKFZp686P01248 antibody, FLJ36156 antibody, HILI 2 antibody, HILI2 antibody, HIWI 2 antibody, HIWI2 antibody, Miwi 2 protein antibody, Miwi2 antibody, PIWI antibody, Piwi like 2 antibody, Piwi like 4 (Drosophila) antibody, Piwi like 4 antibody, Piwi like protein 4 antibody, Piwi like protein antibody, Piwi like RNA mediated gene silencing 4 antibody, Piwi-like protein 4 antibody, PIWIL 4 antibody, PiwiI4 antibody, PIWL4_HUMAN antibody

UniProt:

Q7Z3Z4

Application Details


Application Notes:

Storage Comment:

Buffer: Preservative: 0.03 % Proclin 300 Constituents: 50 % Glycerol, 0.01M PBS, pH 7.4		
Format: Liquid Buffer: Preservative: 0.03 % Proclin 300 Constituents: 50 % Glycerol, 0.01M PBS, pH 7.4 Preservative: ProClin Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.	Restrictions:	For Research Use only
Buffer: Preservative: 0.03 % Proclin 300 Constituents: 50 % Glycerol, 0.01M PBS, pH 7.4 Preservative: ProClin This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.	Handling	
Constituents: 50 % Glycerol, 0.01M PBS, pH 7.4 Preservative: ProClin This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.	Format:	Liquid
Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.	Buffer:	
handled by trained staff only.	Preservative:	ProClin
Storage: -20 °C,-80 °C	Precaution of Use:	
	Storage:	-20 °C,-80 °C


Recommended dilution: WB:1:500-1:5000, IF:1:50-1:200,

Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.



Immunofluorescence

Image 1. Immunofluorescent analysis of A549 cells using ABIN7163478 at dilution of 1:100 and Alexa Fluor 488-congugated AffiniPure Goat Anti-Rabbit IgG(H+L)



Western Blotting

Image 2. Western Blot Positive WB detected in: HEK293 whole cell lysate All lanes: PIWIL4 antibody at 3.4 μg/mL Secondary Goat polyclonal to rabbit IgG at 1/50000 dilution Predicted band size: 97, 57, 53 kDa Observed band size: 97 kDa