antibodies - online.com





Datasheet for ABIN7163603

anti-PAFAH1B1 antibody (AA 40-155) (FITC)



\sim	
()\/\	rview
\cup	1 410 44

Overview	
Quantity:	100 μg
Target:	PAFAH1B1
Binding Specificity:	AA 40-155
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This PAFAH1B1 antibody is conjugated to FITC
Application:	Please inquire
Product Details	
Immunogen:	Recombinant Human Platelet-activating factor acetylhydrolase IB subunit alpha protein (40-
	155AA)
Isotype:	IgG
Cross-Reactivity:	Human
Purification:	>95%, Protein G purified
Target Details	
Target:	PAFAH1B1
Alternative Name:	PAFAH1B1 (PAFAH1B1 Products)
Background:	Background: Required for proper activation of Rho GTPases and actin polymerization at the

leading edge of locomoting cerebellar neurons and postmigratory hippocampal neurons in response to calcium influx triggered via NMDA receptors. Non-catalytic subunit of an acetylhydrolase complex which inactivates platelet-activating factor (PAF) by removing the acetyl group at the SN-2 position (By similarity). Positively regulates the activity of the minusend directed microtubule motor protein dynein. May enhance dynein-mediated microtubule sliding by targeting dynein to the microtubule plus end. Required for several dynein- and microtubule-dependent processes such as the maintenance of Golgi integrity, the peripheral transport of microtubule fragments and the coupling of the nucleus and centrosome. Required during brain development for the proliferation of neuronal precursors and the migration of newly formed neurons from the ventricular/subventricular zone toward the cortical plate. Neuronal migration involves a process called nucleokinesis, whereby migrating cells extend an anterior process into which the nucleus subsequently translocates. During nucleokinesis dynein at the nuclear surface may translocate the nucleus towards the centrosome by exerting force on centrosomal microtubules. May also play a role in other forms of cell locomotion including the migration of fibroblasts during wound healing.

Aliases: LIS 1 antibody, LIS 2 antibody, LIS-1 antibody, LIS1 antibody, LIS1_HUMAN antibody, LIS2 antibody, Lissencephaly 1 protein antibody, Lissencephaly-1 protein antibody, MDCR antibody, MDS antibody, PAF acetylhydrolase 45 kDa subunit antibody, PAF AH 45 kDa subunit antibody, PAF AH alpha antibody, PAF-AH 45 kDa subunit antibody, PAF-AH alpha antibody, PAFAH antibody, PAFAH antibody, PAFAHA antibody, PAFAHA antibody, PAFAHA antibody, PIatelet activating factor acetylhydrolase 1b regulatory subunit 1 antibody, PIatelet activating factor acetylhydrolase IB subunit alpha antibody

UniProt: P43034

Pathways: M Phase, Regulation of Cell Size

Application Details

Restrictions: For Research Use only

Handling

Format: Liquid

Buffer: Preservative: 0.03 % Proclin 300

Constituents: 50 % Glycerol, 0.01M PBS, PH 7.4

Handling

Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C,-80 °C
Storage Comment:	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.