

Datasheet for ABIN7163839

anti-PKD2 antibody (AA 680-968) (HRP)[Go to Product page](#)

Overview

Quantity:	100 µg
Target:	PKD2
Binding Specificity:	AA 680-968
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This PKD2 antibody is conjugated to HRP
Application:	ELISA

Product Details

Immunogen:	Recombinant Human Polycystin-2 protein (680-968AA)
Isotype:	IgG
Cross-Reactivity:	Human
Purification:	>95%, Protein G purified

Target Details

Target:	PKD2
Alternative Name:	PKD2 (PKD2 Products)
Background:	Background: Functions as a cation channel involved in fluid-flow mechanosensation by the primary cilium in renal epithelium (PubMed:18695040). Functions as outward-rectifying K(+)

Target Details

channel, but is also permeable to Ca^{2+} , and to a much lesser degree also to Na^{+} (PubMed:11854751, PubMed:15692563, PubMed:27071085, PubMed:27991905). May contribute to the release of Ca^{2+} stores from the endoplasmic reticulum (PubMed:11854751, PubMed:20881056). Together with TRPV4, forms mechano- and thermosensitive channels in cilium (PubMed:18695040). PKD1 and PKD2 may function through a common signaling pathway that is necessary to maintain the normal, differentiated state of renal tubule cells. Acts as a regulator of cilium length, together with PKD1. The dynamic control of cilium length is essential in the regulation of mechanotransductive signaling. The cilium length response creates a negative feedback loop whereby fluid shear-mediated deflection of the primary cilium, which decreases intracellular cAMP, leads to cilium shortening and thus decreases flow-induced signaling. Also involved in left-right axis specification via its role in sensing nodal flow, forms a complex with PKD1L1 in cilia to facilitate flow detection in left-right patterning. Detection of asymmetric nodal flow gives rise to a Ca^{2+} signal that is required for normal, asymmetric expression of genes involved in the specification of body left-right laterality (By similarity).

Aliases: APKD2 antibody, Autosomal dominant polycystic kidney disease type II antibody, Autosomal dominant polycystic kidney disease type II protein antibody, MGC138466 antibody, MGC138468 antibody, PC 2 antibody, PC2 antibody, PKD 2 antibody, PKD2 antibody, PKD2_HUMAN antibody, PKD4 antibody, Polycystic kidney disease 2 (autosomal dominant) antibody, Polycystic kidney disease 2 antibody, Polycystic kidney disease 2 protein antibody, Polycystin 2 antibody, Polycystin 2 transient receptor potential cation channel antibody, Polycystin-2 antibody, Polycystin2 antibody, Polycystin antibody, R48321 antibody, Transient receptor potential cation channel subfamily P member 2 antibody, TRPP2 antibody

UniProt: [Q13563](#)

Pathways: [cAMP Metabolic Process](#), [Maintenance of Protein Location](#), [Negative Regulation of Transporter Activity](#)

Application Details

Application Notes: Optimal working dilution should be determined by the investigator.

Restrictions: For Research Use only

Handling

Format: Liquid

Handling

Buffer:	Preservative: 0.03 % Proclin 300 Constituents: 50 % Glycerol, 0.01M PBS, pH 7.4
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C,-80 °C
Storage Comment:	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.