
antibodies -online.com





anti-Kcne3 antibody (AA 1-57)

3 Images



Go to Product page

Overview

Quantity:	100 μg
Target:	Kcne3
Binding Specificity:	AA 1-57
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This Kcne3 antibody is un-conjugated
Application:	ELISA, Immunohistochemistry (IHC), Immunofluorescence (IF)

Product Details

Immunogen:	Recombinant Human Potassium voltage-gated channel subfamily E member 3 protein (1-57AA)
Isotype:	IgG
Cross-Reactivity:	Human
Purification:	>95%, Protein G purified

Target Details

Target:	Kcne3
Alternative Name:	KCNE3 (Kcne3 Products)
Background:	Background: Ancillary protein that assembles as a beta subunit with a voltage-gated potassium channel complex of pore-forming alpha subunits. Modulates the gating kinetics and enhances

Target Details

stability of the channel complex. Assembled with KCNB1 modulates the gating characteristics of the delayed rectifier voltage-dependent potassium channel KCNB1 (PubMed:12954870). Associated with KCNC4/Kv3.4 is proposed to form the subthreshold voltage-gated potassium channel in skeletal muscle and to establish the resting membrane potential (RMP) in muscle cells. Associated with KCNQ1/KCLQT1 may form the intestinal cAMP-stimulated potassium channel involved in chloride secretion that produces a current with nearly instantaneous activation with a linear current-voltage relationship.

Aliases: KCNE3 antibody, Potassium voltage-gated channel subfamily E member 3 antibody, MinK-related peptide 2 antibody, Minimum potassium ion channel-related peptide 2 antibody, Potassium channel subunit beta MiRP2 antibody

UniProt:

Q9Y6H6

Application Details

Application Notes:	Recommended dilution: IHC:1:200-1:500, IF:1:50-1:200,
Restrictions:	For Research Use only
Handling	
Format:	Liquid

Buffer: Preservative: 0.03 % Proclin 300

Constituents: 50 % Glycerol, 0.01M PBS, pH 7.4

Preservative:

ProClin


Precaution of Use:

This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Storage:


-20 °C,-80 °C

Storage Comment: Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.




Immunohistochemistry

Image 1. IHC image of ABIN7163983 diluted at 1:200 and staining in paraffin-embedded human cervical cancer performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.



Immunohistochemistry

Image 2. IHC image of ABIN7163983 diluted at 1:200 and staining in paraffin-embedded human glioma performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.



Immunofluorescence

Image 3. Immunofluorescence staining of HepG2 cells with ABIN7163983 at 1:66, counter-stained with DAPI. The cells were fixed in 4% formaldehyde, permeabilized using 0.2% Triton X-100 and blocked in 10% normal Goat Serum. The cells were then incubated with the antibody overnight at 4°C. The secondary antibody was Alexa Fluor 488-congugated AffiniPure Goat Anti-Rabbit IgG(H+L).