Datasheet for ABIN7164090 anti-PRDM16 antibody (AA 448-683) (Biotin)

-online.com antibodies



Overview

| Quantity:            | 100 µg                                       |
|----------------------|--|
| Target:              | PRDM16                                       |
| Binding Specificity: | AA 448-683                                   |
| Reactivity:          | Human  |
| Host:                | Rabbit                                       |
| Clonality:           | Polyclonal                                   |
| Conjugate:           | This PRDM16 antibody is conjugated to Biotin |
| Application:         | ELISA  |

## Product Details

| Immunogen:        | Recombinant Human PR domain zinc finger protein 16 protein (448-683AA) |
|-------------------|--|
| Isotype:          | lgG  |
| Cross-Reactivity: | Human  |
| Purification:     | >95%, Protein G purified   |

## Target Details

| Target:           | PRDM16   |
|-------------------|--|
| Alternative Name: | PRDM16 (PRDM16 Products)   |
| Background:       | Background: Binds DNA and functions as a transcriptional regulator. Functions in the       |
|                   | differentiation of brown adipose tissue (BAT) which is specialized in dissipating chemical |

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/2 | Product datasheet for ABIN7164090 | 09/10/2023 | Copyright antibodies-online. All rights reserved.

Precaution of Use:

Storage Comment:

Storage:

| energy in the form of heat in response to cold or excess feeding while white adipose tissue     |
|---|
| (WAT) is specialized in the storage of excess energy and the control of systemic metabolism.    |
| Together with CEBPB, regulates the differentiation of myoblastic precursors into brown adipose  |
| cells. Functions also as a repressor of TGF-beta signaling. Isoform 4 may regulate granulocytes |
| differentiation.  |
| Aliases: CMD1LL antibody, KIAA1675 antibody, LVNC8 antibody, MDS1/EVI1 like gene 1              |
| antibody, MDS1/EVI1-like gene 1 antibody, MEL1 antibody, PFM 13 antibody, PFM13 antibody,       |
| PR domain containing 16 antibody, PR domain containing protein 16 antibody, PR domain zinc      |
| finger protein 16 antibody, PR domain-containing protein 16 antibody, PRD16_HUMAN antibody,     |
| Prdm16 antibody, Transcription factor MEL 1 antibody, Transcription factor MEL1 antibody        |
| Q9HAZ2  |
| Stem Cell Maintenance, Brown Fat Cell Differentiation   |
|   |
| Optimal working dilution should be determined by the investigator.                              |
| For Research Use only   |
|   |
| Liquid  |
| Preservative: 0.03 % Proclin 300  |
| Constituents: 50 % Glycerol, 0.01M PBS, pH 7.4  |
| ProClin   |
|   |

Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.

handled by trained staff only.

-20 °C,-80 °C

This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be