Datasheet for ABIN7164092 anti-PRDM16 antibody (AA 448-683) (HRP)

antibodies .-online.com



\sim				•			
0	۱	\sim	r\	/1	\sim	۱۸	1
11	\mathbf{V}	-	I \	/ I	-	1/1	
\sim	v	\sim	1 \	/ 1	\sim	v	V

Quantity:	100 µg
Target:	PRDM16
Binding Specificity:	AA 448-683
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This PRDM16 antibody is conjugated to HRP
Application:	ELISA

Product Details

Immunogen:	Recombinant Human PR domain zinc finger protein 16 protein (448-683AA)
Isotype:	lgG
Cross-Reactivity:	Human
Purification:	>95%, Protein G purified

Target Details

Target:	PRDM16
Alternative Name:	PRDM16 (PRDM16 Products)
Background:	Background: Binds DNA and functions as a transcriptional regulator. Functions in the
	differentiation of brown adipose tissue (BAT) which is specialized in dissipating chemical

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/2 | Product datasheet for ABIN7164092 | 09/10/2023 | Copyright antibodies-online. All rights reserved.

Precaution of Use:

Storage Comment:

Storage:

energy in the form of heat in response to cold or excess feeding while white adipose tissue
(WAT) is specialized in the storage of excess energy and the control of systemic metabolism.
Together with CEBPB, regulates the differentiation of myoblastic precursors into brown adipose
cells. Functions also as a repressor of TGF-beta signaling. Isoform 4 may regulate granulocytes
differentiation.
Aliases: CMD1LL antibody, KIAA1675 antibody, LVNC8 antibody, MDS1/EVI1 like gene 1
antibody, MDS1/EVI1-like gene 1 antibody, MEL1 antibody, PFM 13 antibody, PFM13 antibody,
PR domain containing 16 antibody, PR domain containing protein 16 antibody, PR domain zinc
finger protein 16 antibody, PR domain-containing protein 16 antibody, PRD16_HUMAN antibody
Prdm16 antibody, Transcription factor MEL 1 antibody, Transcription factor MEL1 antibody
Q9HAZ2
Stem Cell Maintenance, Brown Fat Cell Differentiation
Optimal working dilution should be determined by the investigator.
For Research Use only
Liquid
Preservative: 0.03 % Proclin 300
Constituents: 50 % Glycerol, 0.01M PBS, pH 7.4
ProClin

Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.

handled by trained staff only.

-20 °C,-80 °C

This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be