antibodies - online.com







anti-DPY19L1 antibody (AA 513-665)



Image



Overview

Quantity:	100 μg
Target:	DPY19L1
Binding Specificity:	AA 513-665
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This DPY19L1 antibody is un-conjugated
Application:	ELISA, Immunofluorescence (IF)

Product Details

Immunogen:	Recombinant Human Probable C-mannosyltransferase DPY19L1 protein (513-665AA)
Isotype:	IgG
Cross-Reactivity:	Human
Purification:	>95%, Protein G purified

Target Details

Target:	DPY19L1
Alternative Name:	DPY19L1 (DPY19L1 Products)
Background:	Background: Probable C-mannosyltransferase that mediates C-mannosylation of tryptophan residues on target proteins.

Target Details

	Aliases: DPY19L1 antibody, GA0500 antibody, KIAA0877Probable C-mannosyltransferase
	DPY19L1 antibody, EC 2.4.1 antibody, Dpy-19-like protein 1 antibody, Protein dpy-19 homolog
	1 antibody
niProt:	Q2PZI1

Uni


Pathways: SARS-CoV-2 Protein Interactome

Application Details

Application Notes:	Recommended dilution: IF:1:50-1:200,
Restrictions:	For Research Use only

Handling	
Format:	Liquid
Buffer:	Preservative: 0.03 % Proclin 300 Constituents: 50 % Glycerol, 0.01M PBS, pH 7.4
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C,-80 °C
Storage Comment:	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.

Images



Immunofluorescence

Image 1. Immunofluorescence staining of A549 cells with ABIN7164389 at 1:166, counter-stained with DAPI. The cells were fixed in 4% formaldehyde, permeabilized using 0.2% Triton X-100 and blocked in 10% normal Goat Serum. The cells were then incubated with the antibody overnight at 4°C. The secondary antibody was Alexa Fluor 488congugated AffiniPure Goat Anti-Rabbit IgG(H+L).