



[Go to Product page](#)

Datasheet for ABIN7164625  
**anti-UTP11 antibody (AA 1-253)**

1 Image

### Overview

Quantity:	100 µL
Target:	UTP11
Binding Specificity:	AA 1-253
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This UTP11 antibody is un-conjugated
Application:	ELISA, Immunohistochemistry (IHC)

### Product Details

Immunogen:	Recombinant Human Probable U3 small nucleolar RNA-associated protein 11 protein (1-253AA)
Isotype:	IgG
Cross-Reactivity:	Human
Purification:	Antigen Affinity Purified

### Target Details

Target:	UTP11
Alternative Name:	UTP11 ( <a href="#">UTP11 Products</a> )
Background:	Background: Involved in nucleolar processing of pre-18S ribosomal RNA.

## Target Details

Aliases: CGI 94 antibody, CGI94 antibody, Probable U3 small nucleolar RNA associated protein 11 antibody, Probable U3 small nucleolar RNA-associated protein 11 antibody, U3 snoRNA associated protein 11 antibody, U3 snoRNA-associated protein 11 antibody, UTP 11L antibody, UTP11 like protein antibody, UTP11 like U3 small nucleolar ribonucleoprotein antibody, UTP11-like protein antibody, UTP11\_HUMAN antibody, Utp11l antibody

UniProt: [Q9Y3A2](#)

## Application Details

Application Notes: Recommended dilution: IHC:1:20-1:200,

Restrictions: For Research Use only

## Handling

Format: Liquid

Buffer: Preservative: 0.03 % Proclin 300  
Constituents: 50 % Glycerol, 0.01M PBS, pH 7.4


Preservative: ProClin

Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Storage: -20 °C,-80 °C

Storage Comment: Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.

## Images



### Immunohistochemistry

**Image 1.** IHC image of ABIN7164625 diluted at 1:100 and staining in paraffin-embedded human brain tissue performed on a Leica Bond™ system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10 % normal goat serum 30 min at RT. Then primary antibody (1 % BSA) was incubated at 4 °C overnight. The primary is detected by a biotinylated secondary antibody and

visualized using an HRP conjugated SP system.