# antibodies -online.com





# anti-PSME4 antibody (AA 105-198) (FITC)



Go to Product page

( )	1/0	r\ /1	014	
( )	ve	I V I	-v	V

Quantity:	100 μg
Target:	PSME4
Binding Specificity:	AA 105-198
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This PSME4 antibody is conjugated to FITC
Application:	Please inquire

### **Product Details**

Immunogen:	Recombinant Human Proteasome activator complex subunit 4 protein (105-198AA)	
Isotype:	IgG	
Cross-Reactivity:	Human	
Purification:	>95%, Protein G purified	

# Target Details

Target:	PSME4	
Alternative Name:	PSME4 (PSME4 Products)	
Background:	Background: Associated component of the proteasome that specifically recognizes acetylate	
	histones and promotes ATP- and ubiquitin-independent degradation of core histones during	

spermatogenesis and DNA damage response. Recognizes and binds acetylated histones via its bromodomain-like (BRDL) region and activates the proteasome by opening the gated channel for substrate entry. Binds to the core proteasome via its C-terminus, which occupies the same binding sites as the proteasomal ATPases, opening the closed structure of the proteasome via an active gating mechanism. Component of the spermatoproteasome, a form of the proteasome specifically found in testis: binds to acetylated histones and promotes degradation of histones, thereby participating actively to the exchange of histones during spermatogenesis. Also involved in DNA damage response in somatic cells, by promoting degradation of histones following DNA double-strand breaks.

Aliases: KIAA0077 antibody, PA200 antibody, Proteasome Activator 200 kDa antibody,
Proteasome activator complex subunit 4 antibody, Proteasome activator PA200 antibody,
PSME 4 antibody, psme4 antibody, PSME4\_HUMAN antibody

UniProt:

Q14997

Pathways:

DNA Replication, Synthesis of DNA

### **Application Details**

Restrictions:

For Research Use only

## Handling

Format:	Liquid	
Buffer:	Preservative: 0.03 % Proclin 300  Constituents: 50 % Glycerol, 0.01M PBS, pH 7.4	
Preservative:	ProClin	
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.	
Storage:	-20 °C,-80 °C	
Storage Comment:	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.	