antibodies .- online.com





anti-PSMB2 antibody (AA 1-201) (FITC)



Overview

| Quantity: | 100 μg |
|----------------------|---|
| Target: | PSMB2 |
| Binding Specificity: | AA 1-201 |
| Reactivity: | Human |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Conjugate: | This PSMB2 antibody is conjugated to FITC |
| Application: | Please inquire |

Product Details

| Immunogen: | Recombinant Human Proteasome subunit beta type-2 protein (1-201AA) |
|-------------------|--|
| Isotype: | IgG |
| Cross-Reactivity: | Human |
| Purification: | >95%, Protein G purified |

Target Details

| Target: | PSMB2 |
|-------------------|--|
| Alternative Name: | PSMB2 (PSMB2 Products) |
| Background: | Background: The proteasome is a multicatalytic proteinase complex with a highly ordered ring |
| | shaped 20S core structure. The core structure is composed of 4 rings of 28 non identical |

subunits, 2 rings are composed of 7 alpha subunits and 2 rings are composed of 7 beta subunits. Proteasomes are distributed throughout eukaryotic cells at a high concentration and cleave peptides in an ATP/ubiquitin dependent process in a non-lysosomal pathway. An essential function of a modified proteasome, the immunoproteasome, is the processing of class I MHC peptides. PSMB2 is a member of the proteasome B-type family, also known as the T1B family, that is a 20S core beta subunit. The proteasome is a multicatalytic proteinase complex with a highly ordered ring shaped 20S core structure. The core structure is composed of 4 rings of 28 non identical subunits, 2 rings are composed of 7 alpha subunits and 2 rings are composed of 7 beta subunits. Proteasomes are distributed throughout eukaryotic cells at a high concentration and cleave peptides in an ATP/ubiquitin dependent process in a non lysosomal pathway. An essential function of a modified proteasome, the immunoproteasome, is the processing of class I MHC peptides. This gene encodes a member of the proteasome B type family, also known as the T1B family, that is a 20S core beta subunit. Publication Note: This RefSeq record includes a subset of the publications that are available for this gene. Please see the Entrez Gene record to access additional publications.

Aliases: HC7 I antibody, Macropain subunit C7 I antibody, Macropain subunit C7-I antibody, Multicatalytic endopeptidase complex subunit C7 1 antibody, Multicatalytic endopeptidase complex subunit C7 I antibody, Multicatalytic endopeptidase complex subunit C7-I antibody, Proteasome (prosome, macropain) subunit beta type 2 antibody, Proteasome beta 2 subunit antibody, Proteasome component C7 I antibody, Proteasome component C7-I antibody, Proteasome subunit beta type-2 antibody, PSB2_HUMAN antibody, Psmb2 antibody

UniProt:

P49721

Pathways:

Mitotic G1-G1/S Phases, DNA Replication, Synthesis of DNA

Application Details

Restrictions:

For Research Use only

Handling

| Format: | Liquid |
|--------------------|--|
| Buffer: | Preservative: 0.03 % Proclin 300 Constituents: 50 % Glycerol, 0.01M PBS, PH 7.4 |
| Preservative: | ProClin |
| Precaution of Use: | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be |

Handling

| | handled by trained staff only. |
|------------------|---|
| Storage: | -20 °C,-80 °C |
| Storage Comment: | Upon receipt, store at -20°C or -80°C. Avoid repeated freeze. |