

[Go to Product page](#)

Datasheet for ABIN7164976

anti-PSMB2 antibody (AA 1-201) (HRP)

Overview

Quantity:	100 µg
Target:	PSMB2
Binding Specificity:	AA 1-201
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This PSMB2 antibody is conjugated to HRP
Application:	ELISA

Product Details

Immunogen:	Recombinant Human Proteasome subunit beta type-2 protein (1-201AA)
Isotype:	IgG
Cross-Reactivity:	Human
Purification:	>95%, Protein G purified

Target Details

Target:	PSMB2
Alternative Name:	PSMB2 (PSMB2 Products)
Background:	Background: The proteasome is a multicatalytic proteinase complex with a highly ordered ring shaped 20S core structure. The core structure is composed of 4 rings of 28 non identical

Target Details

subunits, 2 rings are composed of 7 alpha subunits and 2 rings are composed of 7 beta subunits. Proteasomes are distributed throughout eukaryotic cells at a high concentration and cleave peptides in an ATP/ubiquitin dependent process in a non-lysosomal pathway. An essential function of a modified proteasome, the immunoproteasome, is the processing of class I MHC peptides. PSMB2 is a member of the proteasome B-type family, also known as the T1B family, that is a 20S core beta subunit. The proteasome is a multicatalytic proteinase complex with a highly ordered ring shaped 20S core structure. The core structure is composed of 4 rings of 28 non identical subunits, 2 rings are composed of 7 alpha subunits and 2 rings are composed of 7 beta subunits. Proteasomes are distributed throughout eukaryotic cells at a high concentration and cleave peptides in an ATP/ubiquitin dependent process in a non lysosomal pathway. An essential function of a modified proteasome, the immunoproteasome, is the processing of class I MHC peptides. This gene encodes a member of the proteasome B type family, also known as the T1B family, that is a 20S core beta subunit. Publication Note: This RefSeq record includes a subset of the publications that are available for this gene. Please see the Entrez Gene record to access additional publications.

Aliases: HC7 I antibody, Macropain subunit C7 I antibody, Macropain subunit C7-I antibody, Multicatalytic endopeptidase complex subunit C7 1 antibody, Multicatalytic endopeptidase complex subunit C7 I antibody, Multicatalytic endopeptidase complex subunit C7-I antibody, Proteasome (prosome, macropain) subunit beta type 2 antibody, Proteasome beta 2 subunit antibody, Proteasome component C7 I antibody, Proteasome component C7-I antibody, Proteasome subunit beta type-2 antibody, PSB2_HUMAN antibody, Psmb2 antibody

UniProt: [P49721](#)

Pathways: [Mitotic G1-G1/S Phases](#), [DNA Replication](#), [Synthesis of DNA](#)

Application Details

Application Notes: Optimal working dilution should be determined by the investigator.

Restrictions: For Research Use only

Handling

Format: Liquid

Buffer: Preservative: 0.03 % Proclin 300
Constituents: 50 % Glycerol, 0.01M PBS, PH 7.4

Preservative: ProClin

Handling

Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C,-80 °C
Storage Comment:	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.