



[Go to Product page](#)

Datasheet for ABIN7165036  
**anti-AMBP antibody (AA 20-203)**

2 Images

### Overview

Quantity:	100 µg
Target:	AMBP
Binding Specificity:	AA 20-203
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This AMBP antibody is un-conjugated
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (IHC)

### Product Details

Immunogen:	Recombinant Human Protein AMBP protein (20-203AA)
Isotype:	IgG
Cross-Reactivity:	Human
Purification:	>95%, Protein G purified

### Target Details

Target:	AMBP
Alternative Name:	AMBP ( <a href="#">AMBP Products</a> )
Background:	A1M antibody, Alpha 1 microglobulin/bikunin precursor antibody, Alpha 1 microglycoprotein antibody, Alpha-1 microglycoprotein antibody, Alpha-1-microglobulin antibody, AMBP antibody,

## Target Details

AMBP\_HUMAN antibody, Bikunin antibody, Complex-forming glycoprotein heterogeneous in charge antibody, EDC1 antibody, Growth inhibiting protein 19 antibody, HCP antibody, HI-30 antibody, HI30 antibody, IATIL antibody, ITI antibody, ITI-LC antibody, ITIL antibody, Microglobulin (alpha 1) antibody, Protein AMBP antibody, Protein HC antibody, Trypstatin antibody, Uristatin antibody, Uronic-acid-rich protein antibody, UTI antibody

UniProt: [P02760](#)

## Application Details

Application Notes: Recommended dilution: WB:1:500-1:5000, IHC:1:20-1:200,

Restrictions: For Research Use only

## Handling

Format: Liquid

Buffer: Preservative: 0.03 % Proclin 300  
Constituents: 50 % Glycerol, 0.01M PBS, PH 7.4


Preservative: ProClin

Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Storage: -20 °C,-80 °C


Storage Comment: Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.

## Images



### Immunohistochemistry

**Image 1.** IHC image of ABIN7165036 diluted at 1:150 and staining in paraffin-embedded human liver cancer performed on a Leica Bond™ system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a biotinylated secondary antibody and



visualized using an HRP conjugated SP system.

### Western Blotting

**Image 2.** Western Blot Positive WB detected in: HepG2 whole cell lysate, U251 whole cell lysate All lanes: AMBP antibody at 1:1000 Secondary Goat polyclonal to rabbit IgG at 1/50000 dilution Predicted band size: 39 kDa Observed band size: 39 kDa