

Datasheet for ABIN7165166

anti-C190RF12 antibody (AA 65-104)





Overview

Overview	
Quantity:	100 μg
Target:	C190RF12
Binding Specificity:	AA 65-104
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This C190RF12 antibody is un-conjugated
Application:	ELISA, Immunohistochemistry (IHC), Immunofluorescence (IF)
Product Details	
Immunogen:	Recombinant Human Protein C19orf12 protein (65-104AA)
Isotype:	IgG
Cross-Reactivity:	Human
Purification:	>95%, Protein G purified
Target Details	
Target:	C190RF12
Alternative Name:	C19orf12 (C19ORF12 Products)
Background:	C19orf12 antibody, Chromosome 19 open reading frame 12 antibody, CS012_HUMAN antibody,
	Uncharacterized protein C19orf12 antibody

Target Details

UniProt: Q9NSK7


Application Details

Application Notes:	Recommended dilution: IHC:1:20-1:200, IF:1:50-1:200,
Restrictions:	For Research Use only

Handling


Format:	Liquid
Buffer:	Preservative: 0.03 % Proclin 300 Constituents: 50 % Glycerol, 0.01M PBS, pH 7.4
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C,-80 °C
Storage Comment:	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.

Images




Immunohistochemistry

Image 1. Immunohistochemistry of paraffin-embedded human ovarian cancer using ABIN7165166 at dilution of 1:100



Immunohistochemistry

Image 2. Immunohistochemistry of paraffin-embedded human small intestine tissue using ABIN7165166 at dilution of 1:100



Immunofluorescence

Image 3. Immunofluorescence staining of MCF-7 cells with ABIN7165166 at 1:166, counter-stained with DAPI. The cells were fixed in 4% formaldehyde, permeabilized using 0.2% Triton X-100 and blocked in 10% normal Goat Serum. The cells were then incubated with the antibody overnight at 4°C. The secondary antibody was Alexa Fluor 488-congugated AffiniPure Goat Anti-Rabbit IgG(H+L).