# antibodies .- online.com







## anti-RUNX1T1 antibody (AA 416-525) (Biotin)



( )	1/0	r\ /1	014	
( )	ve	I V I	-v	V

Quantity:	100 μg
Target:	RUNX1T1
Binding Specificity:	AA 416-525
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This RUNX1T1 antibody is conjugated to Biotin
Application:	ELISA

### **Product Details**

Immunogen:	Recombinant Human Protein CBFA2T1 protein (416-525AA)	
Isotype:	IgG	
Cross-Reactivity:	Human	
Purification:	>95%, Protein G purified	

## Target Details

Target:	RUNX1T1	
Alternative Name:	RUNX1T1 (RUNX1T1 Products)	
Background:	Background: Transcriptional corepressor which facilitates transcriptional repression via its association with DNA-binding transcription factors and recruitment of other corepressors and	

histone-modifying enzymes (PubMed:12559562, PubMed:15203199). Can repress the expression of MMP7 in a ZBTB33-dependent manner (PubMed:23251453). Can repress transactivation mediated by TCF12 (PubMed:16803958). Acts as a negative regulator of adipogenesis (By similarity).

Aliases: Acute myelogenous leukemia 1 translocation 1 cyclin D related antibody, Acute myelogenous leukemia 1 translocation 1 protein antibody, AML1 MTG8 antibody, AML1T1 antibody, CBFA2T1 antibody, CDR antibody, Core binding factor runt domain alpha subunit 2 translocated to 1 antibody, Core binding factor runt domain alpha subunit 2 translocated to 1 cyclin D related antibody, Cyclin D related protein antibody, Cyclin-D-related protein antibody, Eight twenty one protein antibody, ETO antibody, ETO protein antibody, MGC2796 antibody, MTG 8 antibody, MTG 8b antibody, MTG8 antibody, MTG8 protein antibody, MTG8\_HUMAN antibody, MTG8b antibody, Myeloid translocation gene on 8q22 antibody, Protein CBFA2T1 antibody, Protein ETO antibody, Protein MTG 8 antibody, Protein MTG8 antibody, Runt related transcription factor 1 translocated to 1 (cyclin D related) antibody, Runt related transcription factor 1 translocated to 1 antibody, Runt related transcription factor 1 translocated to 1 antibody, Runt related transcription factor 1 translocated to 1 antibody, Runt related transcription factor 1 translocated to 1 antibody, Runt related transcription factor 1 translocated to 1 antibody, Runt related transcription factor 1 translocated to 1 antibody, Runt related transcription factor 1 translocated to 1 antibody, Runt related transcription factor 1 translocated to 2 antibody, Runt related transcription factor 1 antibody, Runt related trans

UniProt:

Q06455

For Research Use only

#### **Application Details**

Application Notes:	Optimal working dilution should be determined by the investigator.

## Handling

Restrictions:

Format:	Liquid
Buffer:	Preservative: 0.03 % Proclin 300 Constituents: 50 % Glycerol, 0.01M PBS, PH 7.4
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C,-80 °C

Storage Comment:

Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.