



Datasheet for ABIN7165514
anti-BRINP2 antibody (AA 98-199)



[Go to Product page](#)

2 Images

Overview

Quantity:	100 µL
Target:	BRINP2 (FAM5B)
Binding Specificity:	AA 98-199
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This BRINP2 antibody is un-conjugated
Application:	ELISA, Immunohistochemistry (IHC), Immunofluorescence (IF)

Product Details

Immunogen:	Recombinant Human Protein FAM50B protein (98-199AA)
Isotype:	IgG
Cross-Reactivity:	Human
Purification:	>95%, Protein G purified

Target Details

Target:	BRINP2 (FAM5B)
Alternative Name:	FAM5B (FAM5B Products)
Background:	Background: intercellular bridge, nucleoplasm Aliases: D0H6S2654E antibody, D6S2654E antibody, DNA segment, human D6S2654E antibody,

Target Details

FA50B_HUMAN antibody, FAM50B antibody, Family with sequence similarity 50, member B antibody, Protein FAM50B antibody, Protein XAP 5 like antibody, Protein XAP-5-like antibody, X5L antibody, XAP5 like protein antibody

UniProt: [Q9Y247](#)

Application Details

Application Notes: Recommended dilution: IHC:1:200-1:500, IF:1:50-1:200,

Restrictions: For Research Use only

Handling

Format: Liquid

Buffer: Preservative: 0.03 % Proclin 300
Constituents: 50 % Glycerol, 0.01M PBS, pH 7.4


Preservative: ProClin

Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Storage: -20 °C,-80 °C


Storage Comment: Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.

Images



Immunohistochemistry

Image 1. IHC image of ABIN7165514 diluted at 1:200 and staining in paraffin-embedded human brain tissue performed on a Leica Bond™ system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.



Immunofluorescence

Image 2. Immunofluorescence staining of HeLa cells with ABIN7165514 at 1:66, counter-stained with DAPI. The cells were fixed in 4% formaldehyde, permeabilized using 0.2% Triton X-100 and blocked in 10% normal Goat Serum. The cells were then incubated with the antibody overnight at 4°C. The secondary antibody was Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).