



[Go to Product page](#)

Datasheet for ABIN7165838
anti-NTAN1 antibody (AA 219-310) (Biotin)

Overview

Quantity:	100 µg
Target:	NTAN1
Binding Specificity:	AA 219-310
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This NTAN1 antibody is conjugated to Biotin
Application:	ELISA

Product Details

Immunogen:	Recombinant Human Protein N-terminal asparagine amidohydrolase protein (219-310AA)
Isotype:	IgG
Cross-Reactivity:	Human
Purification:	>95%, Protein G purified

Target Details

Target:	NTAN1
Alternative Name:	NTAN1 (NTAN1 Products)
Background:	Background: Side-chain deamidation of N-terminal asparagine residues to aspartate. Required for the ubiquitin-dependent turnover of intracellular proteins that initiate with Met-Asn. These

Target Details

proteins are acetylated on the retained initiator methionine and can subsequently be modified by the removal of N-acetyl methionine by acylaminoacid hydrolase (AAH). Conversion of the resulting N-terminal asparagine to aspartate by PNAD renders the protein susceptible to arginylation, polyubiquitination and degradation as specified by the N-end rule. This enzyme does not act on substrates with internal or C-terminal asparagines and does not act on glutamine residues in any position, nor on acetylated N-terminal peptidyl Asn.

Aliases: NTAN1 antibody, Protein N-terminal asparagine amidohydrolase antibody, EC 3.5.1.121 antibody, Protein NH2-terminal asparagine amidohydrolase antibody, PNAA antibody, Protein NH2-terminal asparagine deamidase antibody, PNAD antibody, Protein N-terminal Asn amidase antibody, Protein N-terminal asparagine amidase antibody, Protein NTN-amidase antibody

UniProt: [Q96AB6](#)

Application Details

Application Notes: Optimal working dilution should be determined by the investigator.

Restrictions: For Research Use only

Handling

Format: Liquid

Buffer: Preservative: 0.03 % Proclin 300
Constituents: 50 % Glycerol, 0.01M PBS, pH 7.4

Preservative: ProClin

Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Storage: -20 °C,-80 °C

Storage Comment: Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.