antibodies -online.com





anti-POGLUT1 antibody (AA 132-392) (HRP)



Go to Product page

| () | 1/0 | r\ /1 | 014 | |
|-----|-----|-------|-----|---|
| () | ve | I V I | -v | V |
| | | | | |

| Quantity: | 100 μL |
|----------------------|--|
| Target: | POGLUT1 |
| Binding Specificity: | AA 132-392 |
| Reactivity: | Human |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Conjugate: | This POGLUT1 antibody is conjugated to HRP |
| Application: | ELISA |

Product Details

| Immunogen: | Recombinant Human Protein O-glucosyltransferase 1 protein (132-392AA) |
|-------------------|---|
| Isotype: | IgG |
| Cross-Reactivity: | Human |
| Purification: | >95%, Protein G purified |

Target Details

| Target: | POGLUT1 | |
|-------------------|--|--|
| Alternative Name: | POGLUT1 (POGLUT1 Products) | |
| Background: | Background: Dual specificity glycosyltransferase that catalyzes the transfer of glucose and xylose from UDP-glucose and UDP-xylose, respectively, to a serine residue found in the | |

consensus sequence of C-X-S-X-P-C (PubMed:21081508, PubMed:21490058, PubMed:21949356, PubMed:27807076). Specifically targets extracellular EGF repeats of protein such as CRB2, F7, F9 and NOTCH2 (PubMed:21081508, PubMed:21490058, PubMed:21949356, PubMed:27807076). Acts as a positive regulator of Notch signaling by mediating O-glucosylation of Notch, leading to regulate muscle development (PubMed:27807076). Notch glucosylation does not affect Notch ligand binding (PubMed:21490058). Required during early development to promote gastrulation: acts by mediating O-glucosylation of CRB2, which is required for CRB2 localization to the cell membrane (By similarity).

Aliases: C3orf9 antibody, CAP10 like 46 kDa protein antibody, CAP10 like protein, 46 kDa antibody, CAP10-like 46 kDa protein antibody, CLP46 antibody, hCLP46 antibody, hRumi antibody, KDELC family like 1 antibody, KDELCL1 antibody, KTEL (Lys Tyr Glu Leu) containing 1 antibody, KTEL motif containing protein 1 antibody, KTEL motif-containing protein 1 antibody, KTEL1 antibody, KTELC1 antibody, MDS010 antibody, MDSRP antibody, Myelodysplastic syndromes relative protein antibody, O-glucosyltransferase rumi homolog antibody, PGLT1_HUMAN antibody, Poglut1 antibody, protein O glucosyltransferase 1 antibody, Protein O-glucosyltransferase 1 antibody, Rumi antibody, x 010 protein antibody

UniProt: Q8NBL1

Pathways: Notch Signaling

Application Details

| Application Notes: | Optimal working dilution should be determined by the investigator. | |
|--------------------|--|--|
| Restrictions: | For Research Use only | |

Handling

| Hariding | |
|--------------------|---|
| Format: | Liquid |
| Buffer: | Preservative: 0.03 % Proclin 300 Constituents: 50 % Glycerol, 0.01M PBS, pH 7.4 |
| Preservative: | ProClin |
| Precaution of Use: | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only. |

Handling

| Storage: | -20 °C,-80 °C |
|------------------|---|
| Storage Comment: | Upon receipt, store at -20°C or -80°C. Avoid repeated freeze. |