

Datasheet for ABIN716638 anti-DDR1 antibody (pTyr513) (Biotin)



Overview

Quantity:	100 μL
Target:	DDR1
Binding Specificity:	pTyr513
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This DDR1 antibody is conjugated to Biotin
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)),
	Immunohistochemistry (Frozen Sections) (IHC (fro))
Product Details	
Immunogen:	KLH conjugated synthetic phosphopeptide derived from human MCK10 around the
	phosphorylation site of Tyr513
Isotype:	IgG
Cross-Reactivity:	Human
Predicted Reactivity:	Mouse,Rat,Cow,Pig,Horse
Purification:	Purified by Protein A.
Target Details	
Target:	DDR1

Target Details

Alternative Name:	MCK10 (DDD1 Droducto)
Alternative Name:	MCK10 (DDR1 Products)
Background:	Synonyms: MCK10 phospho Y513, p-MCK10 phospho Y513, CAK, CD167a, DDR1CD 167,
	CD167, Cell adhesion kinase, DDR 1, DDR, DDR1, Discoidin domain receptor, Discoidin receptor
	tyrosine kinase isoform a, EDDR 1, Epithelial discoidin domain receptor 1, Epithelial specic
	receptor kinase, Mammarian carcinoma kinase 10, Neuroepithelial tyrosine kinase,
	Neurotrophic tyrosine kinase receptor type 4, NTRK 4, PTK 3, PTK 3A protein tyrosine kinase 3
	Receptor tyrosine kinase NEP, RTK 6, TRK E, Tyrosine kinase receptor E, DDR1_HUMAN.
	Background: Receptor tyrosine kinases (RTKs) play a key role in the communication of cells
	with their microenvironment. These molecules are involved in the regulation of cell growth,
	differentiation and metabolism. The protein encoded by this gene is a RTK that is widely
	expressed in normal and transformed epithelial cells and is activated by various types of
	collagen. This protein belongs to a subfamily of tyrosine kinase receptors with a homology
	region to the Dictyostelium discoideum protein discoidin I in their extracellular domain. Its
	autophosphorylation is achieved by all collagens so far tested (type I to type VI). In situ studies
	and Northern-blot analysis showed that expression of this encoded protein is restricted to
	epithelial cells, particularly in the kidney, lung, gastrointestinal tract, and brain. In addition, this
	protein is significantly over-expressed in several human tumors from breast, ovarian,
	esophageal, and pediatric brain. This gene is located on chromosome 6p21.3 in proximity to
	several HLA class I genes. Alternative splicing of this gene results in multiple transcript variants
Gene ID:	780
UniProt:	Q08345
Pathways:	RTK Signaling, Smooth Muscle Cell Migration
Application Details	
Application Notes:	WB 1:300-5000
	IHC-P 1:200-400
	IHC-F 1:100-500
Restrictions:	For Research Use only
	For Research Use only
Restrictions: Handling Format:	For Research Use only Liquid

Handling

Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C for 12 months.
Expiry Date:	12 months