# antibodies - online.com







## anti-PCDH7 antibody (AA 906-1032)



## Image



$\sim$	
( )\/\Di	view
	VICVV

Quantity:	100 μg
Target:	PCDH7
Binding Specificity:	AA 906-1032
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This PCDH7 antibody is un-conjugated
Application:	ELISA, Immunohistochemistry (IHC)

## **Product Details**

Immunogen:	Recombinant Human Protocadherin-7 protein (906-1032AA)
Isotype:	IgG
Cross-Reactivity:	Human
Purification:	>95%, Protein G purified

## Target Details

Target:	PCDH7
Alternative Name:	PCDH7 (PCDH7 Products)
Background:	BH Pcdh antibody, BH protocadherin antibody, BH-Pcdh antibody, BHPCDH antibody, Brain
	heart protocadherin antibody, Brain-heart protocadherin antibody, PCDH 7 antibody, PCDH7

#### **Target Details**

	antibody, PCDH7_HUMAN antibody, Protocadherin 7 antibody, Protocadherin-7 antibody
UniProt:	060245


### **Application Details**

Application Notes:	Recommended dilution: IHC:1:200-1:500,
Restrictions:	For Research Use only

## Handling

Format:	Liquid
Buffer:	Preservative: 0.03 % Proclin 300 Constituents: 50 % Glycerol, 0.01M PBS, pH 7.4
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C,-80 °C
Storage Comment:	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.

#### **Images**



#### **Immunohistochemistry**

Image 1. IHC image of ABIN7166569 diluted at 1:300 and staining in paraffin-embedded human liver cancer performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.