antibodies -online.com







Image



Go to Product page

Overview

Quantity:	100 μg	
Target:	RPH3AL	
Binding Specificity:	AA 1-315	
Reactivity:	Human	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This RPH3AL antibody is un-conjugated	
Application:	Western Blotting (WB), ELISA	

Product Details

Immunogen:	Recombinant Human Rab effector Noc2 protein (1-315AA)
Isotype:	IgG
Cross-Reactivity:	Human, Rat
Purification:	>95%, Protein G purified

Target Details

Target:	RPH3AL
Alternative Name:	RPH3AL (RPH3AL Products)
Background:	Background: Rab GTPase effector involved in the late steps of regulated exocytosis, both in endocrine and exocrine cells By similarity. Acts as a potential RAB3B effector protein in

e	nith	nel	ial	cel	ls
_			ıuı	OCI	·

Aliases: Gm1753 antibody, No C2 domains protein antibody, Noc2 antibody, Rab effector Noc2 antibody, Rabphilin 3A-like (without C2 domains) antibody, Rabphilin 3A-like antibody, Rabphilin-3A-like protein antibody, RPH3L antibody, RPH3L_HUMAN antibody

UniProt: Q9UNE2

Pathways: Carbohydrate Homeostasis, Regulation of G-Protein Coupled Receptor Protein Signaling


Application Details

Application Notes:	Recommended dilution: WB:1:1000-1:5000,
Restrictions:	For Research Use only

Handling

Format:	Liquid
Buffer:	Preservative: 0.03 % Proclin 300 Constituents: 50 % Glycerol, 0.01M PBS, PH 7.4
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C,-80 °C
Storage Comment:	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.

Images



Western Blotting

Image 1. Western blot All lanes: Rab effector Noc2 antibody at 2 μ g/mL Lane 1: Rat liver tissue Lane 2: K562 whole cell lysate Lane 3: A549 whole cell lysate Secondary Goat polyclonal to rabbit IgG at 1/10000 dilution Predicted band size: 35, 32 kDa Observed band size: 35 kDa