

Datasheet for ABIN7166989

anti-RAB11FIP5 antibody (AA 374-545)**2** Images[Go to Product page](#)

Overview

Quantity:	100 µg
Target:	RAB11FIP5
Binding Specificity:	AA 374-545
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This RAB11FIP5 antibody is un-conjugated
Application:	ELISA, Immunohistochemistry (IHC)

Product Details

Immunogen:	Recombinant Human Rab11 family-interacting protein 5 protein (374-545AA)
Isotype:	IgG
Cross-Reactivity:	Human
Purification:	>95%, Protein G purified

Target Details

Target:	RAB11FIP5
Alternative Name:	RAB11FIP5 (RAB11FIP5 Products)
Background:	Background: Rab effector involved in protein trafficking from apical recycling endosomes to the apical plasma membrane. Involved in insulin granule exocytosis. May regulate V-ATPase

Target Details

intracellular transport in response to extracellular acidosis.

Aliases: GAF 1 antibody, Gaf-1 antibody, GAF1 antibody, gamma SNAP associated factor 1 antibody, Gamma-SNAP-associated factor 1 antibody, KIAA0857 antibody, Phosphoprotein pp75 antibody, pp75 antibody, RAB11 family interacting protein 5 (class I) antibody, Rab11 family-interacting protein 5 antibody, Rab11 FIP5 antibody, Rab11 interacting protein antibody, Rab11 interacting protein Rip11 antibody, Rab11-FIP5 antibody, Rab11-interacting protein Rip11 antibody, Rab11fip5 antibody, RFIP5_HUMAN antibody, RIP11 antibody

UniProt: [Q9BXF6](#)

Pathways: [Hormone Transport](#), [Carbohydrate Homeostasis](#)

Application Details

Application Notes: Recommended dilution: IHC:1:200-1:500,

Restrictions: For Research Use only

Handling

Format: Liquid


Buffer: Preservative: 0.03 % Proclin 300
Constituents: 50 % Glycerol, 0.01M PBS, pH 7.4

Preservative: ProClin

Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.


Storage: -20 °C,-80 °C

Storage Comment: Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.



Immunohistochemistry

Image 1. IHC image of ABIN7166989 diluted at 1:400 and staining in paraffin-embedded human kidney tissue performed on a Leica Bond™ system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.



Immunohistochemistry

Image 2. IHC image of ABIN7166989 diluted at 1:400 and staining in paraffin-embedded human placenta tissue performed on a Leica Bond™ system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.