antibodies - online.com







anti-REST antibody (AA 250-350)



Image



Overview

Quantity:	100 μg
Target:	REST
Binding Specificity:	AA 250-350
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This REST antibody is un-conjugated
Application:	ELISA, Immunofluorescence (IF)

Product Details

Immunogen:	Recombinant Human RE1-silencing transcription factor protein (250-350AA)
Isotype:	IgG
Cross-Reactivity:	Human
Purification:	>95%, Protein G purified

Target Details

Target:	REST
Alternative Name:	REST (REST Products)
Background:	Background: Transcriptional repressor which binds neuron-restrictive silencer element (NRSE)
	and represses neuronal gene transcription in non-neuronal cells. Restricts the expression of

neuronal genes by associating with two distinct corepressors, mSin3 and CoREST, which in turn recruit histone deacetylase to the promoters of REST-regulated genes. Mediates repression by recruiting the BHC complex at RE1/NRSE sites which acts by deacetylating and demethylating specific sites on histones, thereby acting as a chromatin modifier.

Transcriptional repression by REST-CDYL via the recruitment of histone methyltransferase EHMT2 may be important in transformation suppression.

Aliases: Neural Restrictive Silencer Factor antibody, Neural-restrictive silencer factor antibody, Neuron Restrictive Silencer Factor antibody, NRSF antibody, RE1 silencing transcription factor antibody, RE1-silencing transcription factor antibody, Repressor binding to the X2 box antibody, rest antibody, REST protein antibody, REST_HUMAN antibody, X2 box repressor antibody, XBR antibody

UniProt: Q13127

Pathways: Negative Regulation of Hormone Secretion, Regulation of Hormone Metabolic Process,

Regulation of Hormone Biosynthetic Process, Chromatin Binding, Positive Regulation of

Endopeptidase Activity


Application Details

Application Notes: Recommended dilution: IF:1:50-1:200,

Restrictions: For Research Use only

Handling

Format:	Liquid
Buffer:	Preservative: 0.03 % Proclin 300 Constituents: 50 % Glycerol, 0.01M PBS, pH 7.4
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C,-80 °C
Storage Comment:	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.



Immunofluorescence

Image 1. Immunofluorescent analysis of HepG2 cells using ABIN7167425 at dilution of 1:100 and Alexa Fluor 488-congugated AffiniPure Goat Anti-Rabbit IgG(H+L)