

Datasheet for ABIN7167426 anti-REST antibody (AA 250-350) (Biotin)



o to i roudot page

Overview	
Quantity:	100 µg
Target:	REST
Binding Specificity:	AA 250-350
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This REST antibody is conjugated to Biotin
Application:	ELISA
Product Details	
Immunogen:	Recombinant Human RE1-silencing transcription factor protein (250-350AA)
Isotype:	lgG
Cross-Reactivity:	Human

Target Details

Purification:

Target:	REST
Alternative Name:	REST (REST Products)
Background:	Background: Transcriptional repressor which binds neuron-restrictive silencer element (NRSE)
	and represses neuronal gene transcription in non-neuronal cells. Restricts the expression of

>95%, Protein G purified

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/2 | Product datasheet for ABIN7167426 | 07/25/2024 | Copyright antibodies-online. All rights reserved.

	neuronal genes by associating with two distinct corepressors, mSin3 and CoREST, which in
	turn recruit histone deacetylase to the promoters of REST-regulated genes. Mediates
	repression by recruiting the BHC complex at RE1/NRSE sites which acts by deacetylating and
	demethylating specific sites on histones, thereby acting as a chromatin modifier.
	Transcriptional repression by REST-CDYL via the recruitment of histone methyltransferase
	EHMT2 may be important in transformation suppression.
	Aliases: Neural Restrictive Silencer Factor antibody, Neural-restrictive silencer factor antibody,
	Neuron Restrictive Silencer Factor antibody, NRSF antibody, RE1 silencing transcription factor
	antibody, RE1-silencing transcription factor antibody, Repressor binding to the X2 box antibody,
	rest antibody, REST protein antibody, REST_HUMAN antibody, X2 box repressor antibody, XBR
	antibody
UniProt:	Q13127
Pathways:	Negative Regulation of Hormone Secretion, Regulation of Hormone Metabolic Process,
	Regulation of Hormone Biosynthetic Process, Chromatin Binding, Positive Regulation of
	Endopeptidase Activity

Application Details

Application Notes:	Optimal working dilution should be determined by the investigator.
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Buffer:	Preservative: 0.03 % Proclin 300 Constituents: 50 % Glycerol, 0.01M PBS, pH 7.4
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C,-80 °C
Storage Comment:	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.