# antibodies - online.com







## anti-PTPRK antibody (AA 589-749)

**Images** 



( )	11	$\sim$	rv		۱ ۸
	1 \ /	⊢	I \/	╙	1/1

Quantity:	100 μg
Target:	PTPRK
Binding Specificity:	AA 589-749
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This PTPRK antibody is un-conjugated
Application:	Immunohistochemistry (IHC), ELISA, Immunofluorescence (IF)

#### **Product Details**

Immunogen:	Recombinant Human Receptor-type tyrosine-protein phosphatase kappa protein (589-749AA)
Isotype:	IgG
Cross-Reactivity:	Human
Purification:	>95%, Protein G purified

### Target Details

Target:	PTPRK	
Alternative Name:	PTPRK (PTPRK Products)	
Background:	Background: Regulation of processes involving cell contact and adhesion such as growth	
	control, tumor invasion, and metastasis. Negative regulator of EGFR signaling pathway. Forms	

complexes with beta-catenin and gamma-catenin/plakoglobin. Beta-catenin may be a substrate for the catalytic activity of PTPRK/PTP-kappa.

Aliases: dJ480J14.2.1 antibody, DKFZp686C2268 antibody, DKFZp779N1045 antibody, OTTHUMP00000017180 antibody, OTTHUMP00000017181 antibody, OTTHUMP00000017182 antibody, OTTHUMP00000017187 antibody, OTTHUMP00000040306 antibody, Protein tyrosine phosphatase kappa antibody, Protein tyrosine phosphatase kappa precursor antibody, Protein tyrosine phosphatase receptor type K antibody, Protein tyrosine phosphatase receptor type kappa antibody, Protein-tyrosine phosphatase kappa antibody, PTPK antibody, PTPkappa antibody, Ptprk antibody, PTPRK\_HUMAN antibody, R PTP kappa antibody, Receptor type tyrosine protein phosphatase kappa antibody, Receptor-type tyrosine-protein phosphatase kappa antibody

UniProt:

015262

Pathways:

Cell-Cell Junction Organization

#### **Application Details**

Application Notes:


Recommended dilution: IHC:1:200-1:500, IF:1:50-1:200,


Restrictions:

For Research Use only

#### Handling

Format:	Liquid
Buffer:	Preservative: 0.03 % Proclin 300 Constituents: 50 % Glycerol, 0.01M PBS, pH 7.4
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C,-80 °C
Storage Comment:	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.





#### **Immunofluorescence**

**Image 1.** Immunofluorescence staining of HepG2 cells with ABIN7167509 at 1:133, counter-stained with DAPI. The cells were fixed in 4% formaldehyde, permeabilized using 0.2% Triton X-100 and blocked in 10% normal Goat Serum. The cells were then incubated with the antibody overnight at 4°C. The secondary antibody was Alexa Fluor 488-congugated AffiniPure Goat Anti-Rabbit IgG(H+L).

#### **Immunohistochemistry**

Image 2. IHC image of ABIN7167509 diluted at 1:400 and staining in paraffin-embedded human colon cancer performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.

#### **Immunohistochemistry**

Image 3. IHC image of ABIN7167509 diluted at 1:400 and staining in paraffin-embedded human heart tissue performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.