antibodies -online.com







anti-Retinoic Acid Receptor gamma antibody (AA 13-87) (HRP)



| () | 11/0 | K\ / | iew | 1 |
|-----|--------|------|---------------------|---|
| | \cup | ועוי | $\square \vee \vee$ | ı |
| | | | | |

| Quantity: | 100 μg | |
|----------------------|---|--|
| Target: | Retinoic Acid Receptor gamma (RARG) | |
| Binding Specificity: | AA 13-87 | |
| Reactivity: | Human | |
| Host: | Rabbit | |
| Clonality: | Polyclonal | |
| Conjugate: | This Retinoic Acid Receptor gamma antibody is conjugated to HRP | |
| Application: | ELISA | |

Product Details

| Immunogen: | Recombinant Human Retinoic acid receptor gamma protein (13-87AA) | |
|-------------------|--|--|
| Isotype: | IgG | |
| Cross-Reactivity: | Human | |
| Purification: | >95%, Protein G purified | |

Target Details

| Target: | Retinoic Acid Receptor gamma (RARG) |
|-------------------|---|
| Alternative Name: | RARG (RARG Products) |
| Background: | Background: Receptor for retinoic acid. Retinoic acid receptors bind as heterodimers to their |
| | target response elements in response to their ligands, all-trans or 9-cis retinoic acid, and |

regulate gene expression in various biological processes. The RAR/RXR heterodimers bind to the retinoic acid response elements (RARE) composed of tandem 5\'-AGGTCA-3\' sites known as DR1-DR5. In the absence of ligand, acts mainly as an activator of gene expression due to weak binding to corepressors. Required for limb bud development. In concert with RARA or RARB, required for skeletal growth, matrix homeostasis and growth plate function (By similarity).

Aliases: NR1B3 antibody, Nuclear receptor subfamily 1 group B member 3 antibody, RAR gamma antibody, RAR gamma isoform 2 antibody, RAR gamma2 antibody, RAR-gamma antibody, RARC antibody, Rarg antibody, RARG_HUMAN antibody, retinoic acid nuclear receptor gamma variant 1 antibody, retinoic acid nuclear receptor gamma variant 2 antibody, Retinoic acid receptor gamma antibody

UniProt: P13631

Pathways: Nuclear Receptor Transcription Pathway, Retinoic Acid Receptor Signaling Pathway, Steroid

Hormone Mediated Signaling Pathway, Regulation of Cell Size

Application Details

| Application Notes: | Optimal working dilution should be determined by the investigator. |
|--------------------|--|
| | |

Restrictions: For Research Use only

Handling

| Format: | Liquid | |
|--------------------|---|--|
| Buffer: | Preservative: 0.03 % Proclin 300 Constituents: 50 % Glycerol, 0.01M PBS, pH 7.4 | |
| Preservative: | ProClin | |
| Precaution of Use: | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only. | |
| Storage: | -20 °C,-80 °C | |
| Storage Comment: | Upon receipt, store at -20°C or -80°C. Avoid repeated freeze. | |