

Datasheet for ABIN716788 anti-ENPP2 antibody (AA 131-230) (Biotin)



Overview

Quantity:	100 μL
Target:	ENPP2
Binding Specificity:	AA 131-230
Reactivity:	Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This ENPP2 antibody is conjugated to Biotin
Application:	ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human ENPP2
Isotype:	IgG
Cross-Reactivity:	Mouse
Predicted Reactivity:	Human,Rat,Dog,Cow,Pig,Horse,Chicken,Rabbit
Purification:	Purified by Protein A.

Target Details

Target:	ENPP2
Alternative Name:	Enpp2 (ENPP2 Products)

Target Details	
Background:	Synonyms: ATX, Autotaxin,E NPP 2, E-NPP 2, Ectonucleotide
	pyrophosphatase/phosphodiesterase 2, Ectonucleotide pyrophosphatase/phosphodiesterase
	family member 2, Enpp2, ENPP2_HUMAN, Extracellular lysophospholipase D, LysoPLD, PDNP2
	Background: Hydrolyzes lysophospholipids to produce lysophosphatidic acid (LPA) in
	extracellular fluids. Major substrate is lysophosphatidylcholine. Also can act on
	sphingosylphosphphorylcholine producing sphingosine-1-phosphate, a modulator of cell
	motility. Can hydrolyze, in vitro, bis-pNPP, to some extent pNP-TMP, and barely ATP. Involved in
	several motility-related processes such as angiogenesis and neurite outgrowth. Acts as an
	angiogenic factor by stimulating migration of smooth muscle cells and microtubule formation.
	Stimulates migration of melanoma cells, probably via a pertussis toxin-sensitive G protein. May
	have a role in induction of parturition. Possible involvement in cell proliferation and adipose
	tissue development. Tumor cell motility-stimulating factor.
Gene ID:	5168
UniProt:	Q13822
Application Details	
Application Notes:	IHC-P 1:200-400
	IHC-F 1:100-500
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1 μg/μL

Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C for 12 months.
Expiry Date:	12 months