antibodies -online.com





anti-RHOH antibody (AA 1-188)

2 Images



Go to Product page

Overview

| Quantity: | 100 μg |
|----------------------|--|
| Target: | RHOH |
| Binding Specificity: | AA 1-188 |
| Reactivity: | Human |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Conjugate: | This RHOH antibody is un-conjugated |
| Application: | ELISA, Immunofluorescence (IF), Immunohistochemistry (IHC) |

Product Details

| Immunogen: | Recombinant Human Rho-related GTP-binding protein RhoH protein (1-188AA) |
|-------------------|--|
| Isotype: | IgG |
| Cross-Reactivity: | Human |
| Purification: | >95%, Protein G purified |

Target Details

| Target: | RHOH |
|-------------------|---|
| Alternative Name: | RHOH (RHOH Products) |
| Background: | Background: Negative regulator of hematopoietic progenitor cell proliferation, survival and |
| | migration. Critical regulator of thymocyte development and T-cell antigen receptor (TCR) |

signaling by mediating recruitment and activation of ZAP70. Required for phosphorylation of CD3Z, membrane translocation of ZAP70 and subsequent activation of the ZAP70-mediated pathways. Essential for efficient beta-selection and positive selection by promoting the ZAP70dependent phosphorylation of the LAT signalosome during pre-TCR and TCR signaling. Crucial for thymocyte maturation during DN3 to DN4 transition and during positive selection. Plays critical roles in mast cell function by facilitating phosphorylation of SYK in Fc epsilon RImediated signal transduction. Essential for the phosphorylation of LAT, LCP2, PLCG1 and PLCG2 and for Ca(2+) mobilization in mast cells. Binds GTP but lacks intrinsic GTPase activity and is resistant to Rho-specific GTPase-activating proteins. Inhibits the activation of NF-kappa-B by TNF and IKKB and the activation of CRK/p38 by TNF. Inhibits activities of RAC1, RHOA and CDC42. Negatively regulates leukotriene production in neutrophils. Aliases: ARHH antibody, GTP binding protein TTF antibody, GTP-binding protein TTF antibody, OTTHUMP00000125213 antibody, Ras homolog gene family member H antibody, Rho related

GTP binding protein antibody, Rho related GTP binding protein RhoH antibody, Rho-related GTPbinding protein RhoH antibody, RHOH antibody, RHOH_HUMAN antibody, Translocation three four protein antibody, TTF antibody, TTF translocation three four antibody

UniProt:

Q15669

Application Details


Application Notes:

Storage Comment:

| Restrictions: | For Research Use only |
|--------------------|---|
| Handling | |
| Format: | Liquid |
| Buffer: | Preservative: 0.03 % Proclin 300 Constituents: 50 % Glycerol, 0.01M PBS, PH 7.4 |
| Preservative: | ProClin |
| Precaution of Use: | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only. |
| Storage: | -20 °C,-80 °C |


Recommended dilution: IHC:1:20-1:200, IF:1:50-1:200,

Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.



Immunohistochemistry

Image 1. Immunohistochemistry of paraffin-embedded human colon cancer using ABIN7168038 at dilution of 1:100



Immunofluorescence

Image 2. Immunofluorescent analysis of A549 cells using ABIN7168038 at dilution of 1:100 and Alexa Fluor 488-congugated AffiniPure Goat Anti-Rabbit IgG(H+L)