

Datasheet for ABIN7168040 anti-RHOH antibody (AA 1-188) (FITC)



OverviewQuantity:100 µgTarget:RHOHBinding Specificity:AA 1-188Reactivity:HumanHost:RabbitClonality:PolyclonalConjugate:This RHOH antibody is conjugated to FITC

Application: Please inquire

Product Details

| Immunogen: | Recombinant Human Rho-related GTP-binding protein RhoH protein (1-188AA) |
|-------------------|--|
| Isotype: | lgG |
| Cross-Reactivity: | Human |
| Purification: | >95%, Protein G purified |

Target Details

| Target: | RHOH |
|-------------------|--|
| Alternative Name: | RHOH (RHOH Products) |
| Background: | Background: Negative regulator of hematopoietic progenitor cell proliferation, survival and migration. Critical regulator of thymocyte development and T-cell antigen receptor (TCR) |

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/2 | Product datasheet for ABIN7168040 | 07/25/2024 | Copyright antibodies-online. All rights reserved.

| signaling by mediating recruitment and activation of ZAP70. Required for phosphorylation o | of |
|--|------|
| CD3Z, membrane translocation of ZAP70 and subsequent activation of the ZAP70-mediated | d |
| pathways. Essential for efficient beta-selection and positive selection by promoting the ZAP | 70- |
| dependent phosphorylation of the LAT signalosome during pre-TCR and TCR signaling. Cruc | cial |
| for thymocyte maturation during DN3 to DN4 transition and during positive selection. Plays | |
| critical roles in mast cell function by facilitating phosphorylation of SYK in Fc epsilon RI- | |
| mediated signal transduction. Essential for the phosphorylation of LAT, LCP2, PLCG1 and | |
| PLCG2 and for Ca(2+) mobilization in mast cells. Binds GTP but lacks intrinsic GTPase activ | /ity |
| and is resistant to Rho-specific GTPase-activating proteins. Inhibits the activation of NF-kap | ра- |
| B by TNF and IKKB and the activation of CRK/p38 by TNF. Inhibits activities of RAC1, RHOA | and |
| CDC42. Negatively regulates leukotriene production in neutrophils. | |
| Aliases: ARHH antibody, GTP binding protein TTF antibody, GTP-binding protein TTF antibod | dy, |
| OTTHUMP00000125213 antibody, Ras homolog gene family member H antibody, Rho relat | ed |
| GTP binding protein antibody, Rho related GTP binding protein RhoH antibody, Rho-related (| GTP- |
| binding protein RhoH antibody, RHOH antibody, RHOH_HUMAN antibody, Translocation thre | ee |
| four protein antibody, TTF antibody, TTF translocation three four antibody | |

UniProt:

Q15669

Application Details

| Restrictions: | For Research Use only |
|--------------------|---|
| Handling | |
| Format: | Liquid |
| Buffer: | Preservative: 0.03 % Proclin 300 Constituents: 50 % Glycerol, 0.01M PBS, PH 7.4 |
| Preservative: | ProClin |
| Precaution of Use: | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only. |
| Storage: | -20 °C,-80 °C |
| Storage Comment: | Upon receipt, store at -20°C or -80°C. Avoid repeated freeze. |