antibodies -online.com





anti-RPP38 antibody (AA 210-283)





Go to Product page

\sim		
()\/(2rv/	iew
UV	-1 V	IUVV

Quantity:	100 μL
Target:	RPP38
Binding Specificity:	AA 210-283
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This RPP38 antibody is un-conjugated
Application:	ELISA, Immunohistochemistry (IHC)

Product Details

Immunogen:	Recombinant Human Ribonuclease P protein subunit p38 protein (210-283AA)
Isotype:	IgG
Cross-Reactivity:	Human
Purification:	>95%, Protein G purified

Target Details

Target:	RPP38
Alternative Name:	RPP38 (RPP38 Products)
Background:	Background: Component of ribonuclease P, a protein complex that generates mature tRNA
	molecules by cleaving their 5\\\'-ends. RPP38 may associate transiently with RNase P RNA as a

Target Details

factor involved in the transport of H1 RNA to the putative site of its assembly in the cell, the nucleolus.

Aliases: RPP38 antibody, Ribonuclease P protein subunit p38 antibody, RNaseP protein p38 antibody, EC 3.1.26.5 antibody

UniProt:

P78345


Application Details

Application Notes:	Recommended dilution: IHC:1:200-1:500,
Restrictions:	For Research Use only

Handling


Format:	Liquid
Buffer:	Preservative: 0.03 % Proclin 300 Constituents: 50 % Glycerol, 0.01M PBS, pH 7.4
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C,-80 °C
Storage Comment:	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.

Images



Immunohistochemistry

Image 1. IHC image of ABIN7168131 diluted at 1:400 and staining in paraffin-embedded human gastric cancer performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.



Immunohistochemistry

Image 2. IHC image of ABIN7168131 diluted at 1:400 and staining in paraffin-embedded human ovarian cancer performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.