

Datasheet for ABIN7168252  
**anti-RNF219 antibody (AA 257-465)**

## 3 Images

[Go to Product page](#)

## Overview

Quantity:	100 µg
Target:	RNF219
Binding Specificity:	AA 257-465
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (IHC)

## Product Details

Immunogen:	Recombinant Human RING finger protein 219 protein (257-465AA)
Isotype:	IgG
Cross-Reactivity:	Human
Purification:	>95%, Protein G purified

## Target Details

Target:	RNF219
Alternative Name:	RNF219 ( <a href="#">RNF219 Products</a> )
Background:	2610206B13Rik antibody, 2810449K13Rik antibody, AI451544 antibody, DKFZp686A01276 antibody, DKFZp686N15250 antibody, DKFZp686O03173 antibody, FLJ13449 antibody, FLJ25774 antibody, MGC35679 antibody, RGD1564706 antibody, RING finger protein 219 antibody, RING finger protein C13orf7 antibody, RN219_HUMAN antibody, RNF219 antibody

## Target Details

UniProt: [Q5W0B1](#)

## Application Details

Application Notes: Recommended dilution: WB:1:500-1:5000, IHC:1:100-1:300,

Restrictions: For Research Use only

## Handling

Format: Liquid

Buffer: Preservative: 0.03 % Proclin 300  
Constituents: 50 % Glycerol, 0.01M PBS, pH 7.4


Preservative: ProClin

Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Storage: -20 °C,-80 °C


Storage Comment: Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.

## Images




### Immunohistochemistry

**Image 1.** IHC image of ABIN7168252 diluted at 1:300 and staining in paraffin-embedded human melanoma performed on a Leica Bond™ system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10 % normal goat serum 30 min at RT. Then primary antibody (1 % BSA) was incubated at 4 °C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.



### Western Blotting

**Image 2.** Western Blot Positive WB detected in: Hela whole cell lysate All lanes: RNF219 antibody at 4 µg/mL Secondary Goat polyclonal to rabbit IgG at 1/50000 dilution Predicted band size: 81 kDa Observed band size: 81 kDa



### Immunohistochemistry

**Image 3.** IHC image of ABIN7168252 diluted at 1:300 and staining in paraffin-embedded human ovarian cancer performed on a Leica Bond™ system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10 % normal goat serum 30 min at RT. Then primary antibody (1 % BSA) was incubated at 4 °C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.