

[Go to Product page](#)

Datasheet for ABIN7168301

**anti-PAF1/PD2 antibody (AA 1-300) (FITC)**

## Overview

Quantity:	100 µg
Target:	PAF1/PD2 (PAF1)
Binding Specificity:	AA 1-300
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This PAF1/PD2 antibody is conjugated to FITC
Application:	Please inquire

## Product Details

Immunogen:	Recombinant Human RNA polymerase II-associated factor 1 homolog protein (1-300AA)
Isotype:	IgG
Cross-Reactivity:	Human
Purification:	>95%, Protein G purified

## Target Details

Target:	PAF1/PD2 (PAF1)
Alternative Name:	PAF1 ( <a href="#">PAF1 Products</a> )
Background:	Background: Component of the PAF1 complex (PAF1C) which has multiple functions during transcription by RNA polymerase II and is implicated in regulation of development and

## Target Details

maintenance of embryonic stem cell pluripotency. PAF1C associates with RNA polymerase II through interaction with POLR2A CTD non-phosphorylated and 'Ser-2'- and 'Ser-5'-phosphorylated forms and is involved in transcriptional elongation, acting both independently and synergistically with TCEA1 and in cooperation with the DSIF complex and HTATSF1. PAF1C is required for transcription of Hox and Wnt target genes. PAF1C is involved in hematopoiesis and stimulates transcriptional activity of KMT2A/MLL1, it promotes leukemogenesis through association with KMT2A/MLL1-rearranged oncoproteins, such as KMT2A/MLL1-MLLT3/AF9 and KMT2A/MLL1-MLLT1/ENL. PAF1C is involved in histone modifications such as ubiquitination of histone H2B and methylation on histone H3 'Lys-4' (H3K4me3). PAF1C recruits the RNF20/40 E3 ubiquitin-protein ligase complex and the E2 enzyme UBE2A or UBE2B to chromatin which mediate monoubiquitination of 'Lys-120' of histone H2B (H2BK120ub1), UB2A/B-mediated H2B ubiquitination is proposed to be coupled to transcription. PAF1C is involved in mRNA 3' end formation probably through association with cleavage and poly(A) factors. In case of infection by influenza A strain H3N2, PAF1C associates with viral NS1 protein, thereby regulating gene transcription. Connects PAF1C with the RNF20/40 E3 ubiquitin-protein ligase complex. Involved in polyadenylation of mRNA precursors. Has oncogenic activity in vivo and in vitro.

Aliases: F23149 1 antibody, FLJ11123 antibody, hPAF1 antibody, PAF 1 antibody, PAF1 antibody, Paf1 RNA polymerase II associated factor homolog antibody, PAF1\_HUMAN antibody, Pancreatic differentiation protein 2 antibody, PD 2 antibody, PD2 antibody, PD2 protein antibody, RNA polymerase II associated factor homolog antibody, RNA polymerase II-associated factor 1 homolog antibody

UniProt: [Q8N7H5](#)

Pathways: [Cellular Response to Molecule of Bacterial Origin, Stem Cell Maintenance](#)

## Application Details

Restrictions: For Research Use only

## Handling

Format: Liquid

Buffer: Preservative: 0.03 % Proclin 300  
Constituents: 50 % Glycerol, 0.01M PBS, PH 7.4

Preservative: ProClin

## Handling

---

Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C,-80 °C
Storage Comment:	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.