antibodies .- online.com





anti-SCARA5 antibody (AA 303-428) (Biotin)



Go to Product page

\sim			
	N/P	r\/I	i⊢₩

Quantity:	100 μg
Target:	SCARA5
Binding Specificity:	AA 303-428
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This SCARA5 antibody is conjugated to Biotin
Application:	ELISA

Product Details

Immunogen:	Recombinant Human Scavenger receptor class A member 5 protein (303-428AA)	
Isotype:	IgG	
Cross-Reactivity:	Human	
Purification:	>95%, Protein G purified	

Target Details

Target:	SCARA5	
Alternative Name:	SCARA5 (SCARA5 Products)	
Background:	Background: Ferritin receptor that mediates non-transferrin-dependent delivery of iron.	
	Mediates cellular uptake of ferritin-bound iron by stimulating ferritin endocytosis from the cell	

surface with consequent iron delivery within the cell. Delivery of iron to cells by ferritin is required for the development of specific cell types, suggesting the existence of cell typespecific mechanisms of iron traffic in organogenesis, which alternatively utilize transferrin or non-transferrin iron delivery pathways. Ferritin mediates iron uptake in capsule cells of the developing kidney. Binds preferrentially ferritin light chain (FTL) compared to heavy chain (FTH1).

Aliases: FLJ23907 antibody, MGC45780 antibody, NET33 antibody, SCAR5_HUMAN antibody, SCARA 5 antibody, scara5 antibody, Scavenger receptor class A member 5 antibody, Scavenger receptor class A, member 5 (putative) antibody, Scavenger receptor class A, member 5 antibody, Scavenger receptor hlg antibody, Test antibody, Testis expressed scavenger receptor antibody

UniProt: Q6ZMJ2

Pathways: Transition Metal Ion Homeostasis

Application Details

Application Notes: Optimal working dilution should be determined by the investigator.

For Research Use only

Handling

Restrictions:

Format:

Liquid

Buffer:

Preservative: 0.03 % Proclin 300

Constituents: 50 % Glycerol, 0.01M PBS, pH 7.4

Preservative:

ProClin

Precaution of Use:

This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Storage:

-20 °C,-80 °C

Upon receipt, store at -20 °C or -80 °C. Avoid repeated freeze.