

Datasheet for ABIN7169028

anti-CHEK2 antibody (AA 1-270) (Biotin)[Go to Product page](#)

Overview

Quantity:	100 µg
Target:	CHEK2
Binding Specificity:	AA 1-270
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This CHEK2 antibody is conjugated to Biotin
Application:	ELISA

Product Details

Immunogen:	Recombinant Human Serine/threonine-protein kinase Chk2 protein (1-270AA)
Isotype:	IgG
Cross-Reactivity:	Human
Purification:	>95%, Protein G purified

Target Details

Target:	CHEK2
Alternative Name:	CHEK2 (CHEK2 Products)
Background:	Background: Serine/threonine-protein kinase which is required for checkpoint-mediated cell cycle arrest, activation of DNA repair and apoptosis in response to the presence of DNA double-

Target Details

strand breaks. May also negatively regulate cell cycle progression during unperturbed cell cycles. Following activation, phosphorylates numerous effectors preferentially at the consensus sequence [L-X-R-X-X-S/T]. Regulates cell cycle checkpoint arrest through phosphorylation of CDC25A, CDC25B and CDC25C, inhibiting their activity. Inhibition of CDC25 phosphatase activity leads to increased inhibitory tyrosine phosphorylation of CDK-cyclin complexes and blocks cell cycle progression. May also phosphorylate NEK6 which is involved in G2/M cell cycle arrest. Regulates DNA repair through phosphorylation of BRCA2, enhancing the association of RAD51 with chromatin which promotes DNA repair by homologous recombination. Also stimulates the transcription of genes involved in DNA repair (including BRCA2) through the phosphorylation and activation of the transcription factor FOXM1. Regulates apoptosis through the phosphorylation of p53/TP53, MDM4 and PML. Phosphorylation of p53/TP53 at '\Ser-20\' by CHEK2 may alleviate inhibition by MDM2, leading to accumulation of active p53/TP53. Phosphorylation of MDM4 may also reduce degradation of p53/TP53. Also controls the transcription of pro-apoptotic genes through phosphorylation of the transcription factor E2F1. Tumor suppressor, it may also have a DNA damage-independent function in mitotic spindle assembly by phosphorylating BRCA1. Its absence may be a cause of the chromosomal instability observed in some cancer cells. Promotes the CCAR2-SIRT1 association and is required for CCAR2-mediated SIRT1 inhibition (PubMed:25361978).

Aliases: CDS 1 antibody, Cds1 antibody, Cds1 homolog antibody, Checkpoint kinase 2 antibody, Checkpoint like protein CHK2 antibody, CHEK 2 antibody, Chk2 antibody, Chk 2 antibody, CHK2 checkpoint homolog (S. pombe) antibody, CHK2 checkpoint homolog antibody, CHK2_HUMAN antibody, hCds1 antibody, HuCds 1 antibody, LFS 2 antibody, LFS2 antibody, PP1425 antibody, RAD 53 antibody, RAD53 antibody, Rad53 homolog antibody, Serine/threonine protein kinase Chk2 antibody, Serine/threonine-protein kinase Chk2 antibody

UniProt:	O96017
Pathways:	p53 Signaling , Apoptosis , Cell Division Cycle

Application Details

Application Notes:	Optimal working dilution should be determined by the investigator.
Restrictions:	For Research Use only

Handling

Format:	Liquid
---------	--------

Handling

Buffer:	Preservative: 0.03 % Proclin 300 Constituents: 50 % Glycerol, 0.01M PBS, PH 7.4
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C,-80 °C
Storage Comment:	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.