Datasheet for ABIN7169045 anti-MASTL antibody (AA 612-740) (FITC)

-online.com antibodies



Overview

Quantity:	100 µg
Target:	MASTL
Binding Specificity:	AA 612-740
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This MASTL antibody is conjugated to FITC
Application:	Please inquire

## Product Details

Immunogen:	Recombinant Human Serine/threonine-protein kinase greatwall protein (612-740AA)
Isotype:	lgG
Cross-Reactivity:	Human
Purification:	>95%, Protein G purified

## Target Details

Target:	MASTL
Alternative Name:	MASTL (MASTL Products)
Background:	Background: Serine/threonine kinase that plays a key role in M phase by acting as a regulator of
	mitosis entry and maintenance. Acts by promoting the inactivation of protein phosphatase 2A

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/2 | Product datasheet for ABIN7169045 | 09/10/2023 | Copyright antibodies-online. All rights reserved.

5	
Application Details	
UniProt:	Q96GX5
	antibody, THC2 antibody
	protein kinase-like antibody, RP11 85G18.2 antibody, Serine/threonine-protein kinase greatwall
	associated serine/threonine kinase like antibody, Microtubule-associated serine/threonine-
	antibody, hGWL antibody, MAST-L antibody, Mastl antibody, MGC117975 antibody, Microtubule
	antibody, Greatwall protein kinase antibody, GW antibody, GWL antibody, GWL_HUMAN
	Aliases: 2700091H24Rik antibody, C88295 antibody, FLJ14813 antibody, GREATWALL
	be involved in megakaryocyte differentiation.
	inhibited. Phosphorylates histone protein in vitro, however such activity is unsure in vivo. May
	CDK1 activity high. Following DNA damage, it is also involved in checkpoint recovery by being
	delta) subunit of PP2A. Inactivation of PP2A during M phase is essential to keep cyclin-B1-
	ARPP19 and ENSA are phosphatase inhibitors that specifically inhibit the PPP2R2D (PR55-
	and subsequent activation of ARPP19 and ENSA at \'Ser-62\' and \'Ser-67\', respectively.
	(PP2A) during M phase: does not directly inhibit PP2A but acts by mediating phosphorylation

## Restrictions: For Research Use only Handling Format: Liquid Buffer: Preservative: 0.03 % Proclin 300 Constituents: 50 % Glycerol, 0.01M PBS, pH 7.4 Preservative: ProClin Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only. -20 °C,-80 °C Storage: Storage Comment: Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.