

Datasheet for ABIN7169068
anti-MARK2 antibody (AA 373-594)[Go to Product page](#)

2 Images

Overview

Quantity:	100 µg
Target:	MARK2
Binding Specificity:	AA 373-594
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This MARK2 antibody is un-conjugated
Application:	ELISA, Western Blotting (WB), Immunohistochemistry (IHC)

Product Details

Immunogen:	Recombinant Human Serine/threonine-protein kinase MARK2 protein (373-594AA)
Isotype:	IgG
Cross-Reactivity:	Human
Purification:	>95%, Protein G purified

Target Details

Target:	MARK2
Alternative Name:	MARK2 (MARK2 Products)
Background:	Background: Serine/threonine-protein kinase involved in cell polarity and microtubule dynamics regulation. Phosphorylates CRTC2/TORC2, DCX, HDAC7, KIF13B, MAP2, MAP4, MAPT/TAU,

Target Details

and RAB11FIP2. Plays a key role in cell polarity by phosphorylating the microtubule-associated proteins MAP2, MAP4 and MAPT/TAU at KXGS motifs, causing detachment from microtubules, and their disassembly. Regulates epithelial cell polarity by phosphorylating RAB11FIP2.

Involved in the regulation of neuronal migration through its dual activities in regulating cellular polarity and microtubule dynamics, possibly by phosphorylating and regulating DCX. Regulates axogenesis by phosphorylating KIF13B, promoting interaction between KIF13B and 14-3-3 and inhibiting microtubule-dependent accumulation of KIF13B. Also required for neurite outgrowth and establishment of neuronal polarity. Regulates localization and activity of some histone deacetylases by mediating phosphorylation of HDAC7, promoting subsequent interaction between HDAC7 and 14-3-3 and export from the nucleus. Also acts as a positive regulator of the Wnt signaling pathway, probably by mediating phosphorylation of dishevelled proteins (DVL1, DVL2 and/or DVL3). Modulates the developmental decision to build a columnar versus a hepatic epithelial cell apparently by promoting a switch from a direct to a transcytotic mode of apical protein delivery. Essential for the asymmetric development of membrane domains of polarized epithelial cells.

Aliases: ELKL motif kinase 1 antibody, ELKL motif kinase antibody, EMK-1 antibody, EMK1 antibody, MAP/microtubule affinity regulating kinase 2 antibody, MAP/microtubule affinity-regulating kinase 2 antibody, Mark2 antibody, MARK2_HUMAN antibody, MGC99619 antibody, PAR 1 antibody, Par 1b antibody, PAR1 homolog antibody, Par1b antibody, Ser/Thr protein kinase PAR 1B antibody, Serine/threonine protein kinase EMK antibody, Serine/threonine protein kinase MARK2 antibody, Serine/threonine-protein kinase MARK2 antibody

UniProt: [Q7KZI7](#)

Pathways: [SARS-CoV-2 Protein Interactome](#), [The Global Phosphorylation Landscape of SARS-CoV-2 Infection](#)

Application Details

Application Notes: Recommended dilution: WB:1:2000-1:10000, IHC:1:20-1:200,

Restrictions: For Research Use only

Handling


Format: Liquid

Buffer: Preservative: 0.03 % Proclin 300
Constituents: 50 % Glycerol, 0.01M PBS, pH 7.4

Handling


Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C,-80 °C
Storage Comment:	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.

Images



Immunohistochemistry

Image 1. Immunohistochemistry of paraffin-embedded human kidney tissue using ABIN7169068 at dilution of 1:100



Western Blotting

Image 2. Western Blot Positive WB detected in: Jurkat whole cell lysate, K562 whole cell lysate All lanes: MARK2 antibody at 1:2000 Secondary Goat polyclonal to rabbit IgG at 1/50000 dilution Predicted band size: 88, 79, 84, 82, 81, 78, 87, 83, 80 kDa Observed band size: 88 kDa